

Discrete Math for Computer Science Students

Ken Bogart
Dept. of Mathematics
Dartmouth College

Scot Drysdale
Dept. of Computer Science
Dartmouth College

Cliff Stein
Dept. of Industrial Engineering
and Operations Research
Columbia University

Contents

1	Counting	1
1.1	Basic Counting	1
	The Sum Principle	1
	Abstraction	2
	Summing Consecutive Integers	3
	The Product Principle	3
	Two element subsets	5
	Important Concepts, Formulas, and Theorems	6
	Problems	7
1.2	Counting Lists, Permutations, and Subsets.	9
	Using the Sum and Product Principles	9
	Lists and functions	10
	The Bijection Principle	12
	k -element permutations of a set	13
	Counting subsets of a set	13
	Important Concepts, Formulas, and Theorems	15
	Problems	16
1.3	Binomial Coefficients	19
	Pascal's Triangle	19
	A proof using the Sum Principle	20
	The Binomial Theorem	22
	Labeling and trinomial coefficients	23
	Important Concepts, Formulas, and Theorems	24
	Problems	25
1.4	Equivalence Relations and Counting (Optional)	27
	The Symmetry Principle	27

Equivalence Relations	28
The Quotient Principle	29
Equivalence class counting	30
Multisets	31
The bookcase arrangement problem.	32
The number of k -element multisets of an n -element set.	33
Using the quotient principle to explain a quotient	34
Important Concepts, Formulas, and Theorems	34
Problems	35
2 Cryptography and Number Theory	39
2.1 Cryptography and Modular Arithmetic	39
Introduction to Cryptography	39
Private Key Cryptography	40
Public-key Cryptosystems	42
Arithmetic modulo n	43
Cryptography using multiplication mod n	47
Important Concepts, Formulas, and Theorems	48
Problems	49
2.2 Inverses and GCDs	52
Solutions to Equations and Inverses mod n	52
Inverses mod n	53
Converting Modular Equations to Normal Equations	55
Greatest Common Divisors (GCD)	55
Euclid's Division Theorem	56
The GCD Algorithm	58
Extended GCD algorithm	59
Computing Inverses	61
Important Concepts, Formulas, and Theorems	62
Problems	63
2.3 The RSA Cryptosystem	66
Exponentiation mod n	66
The Rules of Exponents	66
Fermat's Little Theorem	68

The RSA Cryptosystem	69
The Chinese Remainder Theorem	72
Important Concepts, Formulas, and Theorems	73
Problems	74
2.4 Details of the RSA Cryptosystem	76
Practical Aspects of Exponentiation mod n	76
How long does it take to use the RSA Algorithm?	77
How hard is factoring?	78
Finding large primes	78
Important Concepts, Formulas, and Theorems	81
Problems	81
3 Reflections on Logic and Proof	83
3.1 Equivalence and Implication	83
Equivalence of statements	83
Truth tables	85
DeMorgan's Laws	88
Implication	89
Important Concepts, Formulas, and Theorems	92
Problems	94
3.2 Variables and Quantifiers	96
Variables and universes	96
Quantifiers	97
Standard notation for quantification	98
Statements about variables	99
Rewriting statements to encompass larger universes	100
Proving quantified statements true or false	101
Negation of quantified statements	101
Implicit quantification	103
Proof of quantified statements	104
Important Concepts, Formulas, and Theorems	105
Problems	106
3.3 Inference	108
Direct Inference (Modus Ponens) and Proofs	108

Rules of inference for direct proofs	109
Contrapositive rule of inference.	110
Proof by contradiction	112
Important Concepts, Formulas, and Theorems	114
Problems	115
4 Induction, Recursion, and Recurrences	117
4.1 Mathematical Induction	117
Smallest Counter-Examples	117
The Principle of Mathematical Induction	120
Strong Induction	123
Induction in general	124
Important Concepts, Formulas, and Theorems	125
Problems	126
4.2 Recursion, Recurrences and Induction	128
Recursion	128
First order linear recurrences	129
Iterating a recurrence	130
Geometric series	131
First order linear recurrences	133
Important Concepts, Formulas, and Theorems	136
Problems	137
4.3 Growth Rates of Solutions to Recurrences	139
Divide and Conquer Algorithms	139
Recursion Trees	140
Three Different Behaviors	146
Important Concepts, Formulas, and Theorems	148
Problems	148
4.4 The Master Theorem	150
Master Theorem	150
Solving More General Kinds of Recurrences	152
More realistic recurrences (Optional)	154
Recurrences for general n (Optional)	155
Appendix: Proofs of Theorems (Optional)	157

	Important Concepts, Formulas, and Theorems	159
	Problems	161
4.5	More general kinds of recurrences	163
	Recurrence Inequalities	163
	A Wrinkle with Induction	164
	Further Wrinkles in Induction Proofs	165
	Dealing with Functions Other Than n^c	167
	Important Concepts, Formulas, and Theorems	171
	Problems	172
4.6	Recurrences and Selection	174
	The idea of selection	174
	A recursive selection algorithm	174
	Selection without knowing the median in advance	175
	An algorithm to find an element in the middle half	177
	An analysis of the revised selection algorithm	179
	Uneven Divisions	180
	Important Concepts, Formulas, and Theorems	182
	Problems	182
5	Probability	185
5.1	Introduction to Probability	185
	Why do we study probability?	185
	Some examples of probability computations	186
	Complementary probabilities	187
	Probability and hashing	188
	The Uniform Probability Distribution	188
	Important Concepts, Formulas, and Theorems	191
	Problems	192
5.2	Unions and Intersections	194
	The probability of a union of events	194
	Principle of inclusion and exclusion for probability	196
	The principle of inclusion and exclusion for counting	200
	Important Concepts, Formulas, and Theorems	201
	Problems	202

5.3	Conditional Probability and Independence	204
	Conditional Probability	204
	Independence	206
	Independent Trials Processes	208
	Tree diagrams	209
	Important Concepts, Formulas, and Theorems	212
	Problems	213
5.4	Random Variables	215
	What are Random Variables?	215
	Binomial Probabilities	215
	Expected Value	218
	Expected Values of Sums and Numerical Multiples	220
	The Number of Trials until the First Success	222
	Important Concepts, Formulas, and Theorems	224
	Problems	225
5.5	Probability Calculations in Hashing	227
	Expected Number of Items per Location	227
	Expected Number of Empty Locations	228
	Expected Number of Collisions	228
	Expected maximum number of elements in a slot of a hash table (Optional)	230
	Important Concepts, Formulas, and Theorems	234
	Problems	235
5.6	Conditional Expectations, Recurrences and Algorithms	237
	When Running Times Depend on more than Size of Inputs	237
	Conditional Expected Values	238
	Randomized algorithms	240
	A more exact analysis of RandomSelect	245
	Important Concepts, Formulas, and Theorems	247
	Problems	248
5.7	Probability Distributions and Variance	251
	Distributions of random variables	251
	Variance	253
	Important Concepts, Formulas, and Theorems	258
	Problems	259

6	Graphs	263
6.1	Graphs	263
	The degree of a vertex	265
	Connectivity	267
	Cycles	269
	Trees	269
	Other Properties of Trees	270
	Important Concepts, Formulas, and Theorems	272
	Problems	274
6.2	Spanning Trees and Rooted Trees	276
	Spanning Trees	276
	Breadth First Search	278
	Rooted Trees	281
	Important Concepts, Formulas, and Theorems	283
	Problems	285
6.3	Eulerian and Hamiltonian Paths and Tours	288
	Eulerian Tours and Trails	288
	Hamiltonian Paths and Cycles	291
	NP -Complete Problems	295
	Important Concepts, Formulas, and Theorems	297
	Problems	298
6.4	Matching Theory	300
	The idea of a matching	300
	Making matchings bigger	303
	Matching in Bipartite Graphs	305
	Searching for Augmenting Paths in Bipartite Graphs	306
	The Augmentation-Cover algorithm	307
	Good Algorithms	309
	Important Concepts, Formulas, and Theorems	310
	Problems	311
6.5	Coloring and planarity	313
	The idea of coloring	313
	Interval Graphs	315
	Planarity	317

The Faces of a Planar Drawing	318
The Five Color Theorem	321
Important Concepts, Formulas, and Theorems	323
Problems	324

Chapter 1

Counting

1.1 Basic Counting

The Sum Principle

We begin with an example that illustrates a fundamental principle.

Exercise 1.1-1 The loop below is part of an implementation of selection sort, which sorts a list of items chosen from an ordered set (numbers, alphabet characters, words, etc.) into non-decreasing order.

```
(1) for  $i = 1$  to  $n - 1$ 
(2)     for  $j = i + 1$  to  $n$ 
(3)         if ( $A[i] > A[j]$ )
(4)             exchange  $A[i]$  and  $A[j]$ 
```

How many times is the comparison $A[i] > A[j]$ made in Line 3?

In Exercise 1.1-1, the segment of code from lines 2 through 4 is executed $n - 1$ times, once for each value of i between 1 and $n - 1$ inclusive. The first time, it makes $n - 1$ comparisons. The second time, it makes $n - 2$ comparisons. The i th time, it makes $n - i$ comparisons. Thus the total number of comparisons is

$$(n - 1) + (n - 2) + \cdots + 1 . \tag{1.1}$$

This formula is not as important as the reasoning that lead us to it. In order to put the reasoning into a broadly applicable format, we will describe what we were doing in the language of sets. Think about the set S containing all comparisons the algorithm in Exercise 1.1-1 makes. We divided set S into $n - 1$ pieces (i.e. smaller sets), the set S_1 of comparisons made when $i = 1$, the set S_2 of comparisons made when $i = 2$, and so on through the set S_{n-1} of comparisons made when $i = n - 1$. We were able to figure out the number of comparisons in each of these pieces by observation, and added together the sizes of all the pieces in order to get the size of the set of all comparisons.

in order to describe a general version of the process we used, we introduce some set-theoretic terminology. Two sets are called *disjoint* when they have no elements in common. Each of the sets S_i we described above is disjoint from each of the others, because the comparisons we make for one value of i are different from those we make with another value of i . We say the set of sets $\{S_1, \dots, S_m\}$ (above, m was $n - 1$) is a family of *mutually disjoint sets*, meaning that it is a family (set) of sets, any two of which are disjoint. With this language, we can state a general principle that explains what we were doing without making any specific reference to the problem we were solving.

Principle 1.1 (Sum Principle) *The size of a union of a family of mutually disjoint finite sets is the sum of the sizes of the sets.*

Thus we were, in effect, using the sum principle to solve Exercise 1.1-1. We can describe the sum principle using an algebraic notation. Let $|S|$ denote the size of the set S . For example, $|\{a, b, c\}| = 3$ and $|\{a, b, a\}| = 2$.¹ Using this notation, we can state the sum principle as: if S_1, S_2, \dots, S_m are disjoint sets, then

$$|S_1 \cup S_2 \cup \dots \cup S_m| = |S_1| + |S_2| + \dots + |S_m|. \quad (1.2)$$

To write this without the “dots” that indicate left-out material, we write

$$|\bigcup_{i=1}^m S_i| = \sum_{i=1}^m |S_i|.$$

When we can write a set S as a union of disjoint sets S_1, S_2, \dots, S_k we say that we have *partitioned* S into the sets S_1, S_2, \dots, S_k , and we say that the sets S_1, S_2, \dots, S_k form a *partition* of S . Thus $\{\{1\}, \{3, 5\}, \{2, 4\}\}$ is a partition of the set $\{1, 2, 3, 4, 5\}$ and the set $\{1, 2, 3, 4, 5\}$ can be partitioned into the sets $\{1\}, \{3, 5\}, \{2, 4\}$. It is clumsy to say we are partitioning a set into sets, so instead we call the sets S_i into which we partition a set S the *blocks* of the partition. Thus the sets $\{1\}, \{3, 5\}, \{2, 4\}$ are the blocks of a partition of $\{1, 2, 3, 4, 5\}$. In this language, we can restate the sum principle as follows.

Principle 1.2 (Sum Principle) *If a finite set S has been partitioned into blocks, then the size of S is the sum of the sizes of the blocks.*

Abstraction

The process of figuring out a general principle that explains why a certain computation makes sense is an example of the mathematical process of *abstraction*. We won't try to give a precise definition of abstraction but rather point out examples of the process as we proceed. In a course in set theory, we would further abstract our work and derive the sum principle from the axioms of

¹It may look strange to have $|\{a, b, a\}| = 2$, but an element either is or is not in a set. It cannot be in a set multiple times. (This situation leads to the idea of multisets that will be introduced later on in this section.) We gave this example to emphasize that the notation $\{a, b, a\}$ means the same thing as $\{a, b\}$. Why would someone even contemplate the notation $\{a, b, a\}$. Suppose we wrote $S = \{x|x \text{ is the first letter of Ann, Bob, or Alice}\}$. Explicitly following this description of S would lead us to first write down $\{a, b, a\}$ and then realize it equals $\{a, b\}$.

set theory. In a course in discrete mathematics, this level of abstraction is unnecessary, so we will simply use the sum principle as the basis of computations when it is convenient to do so. If our goal were only to solve this one exercise, then our abstraction would have been almost a mindless exercise that complicated what was an “obvious” solution to Exercise 1.1-1. However the sum principle will prove to be useful in a wide variety of problems. Thus we observe the value of abstraction—when you can recognize the abstract elements of a problem, then abstraction often helps you solve subsequent problems as well.

Summing Consecutive Integers

Returning to the problem in Exercise 1.1-1, it would be nice to find a simpler form for the sum given in Equation 1.1. We may also write this sum as

$$\sum_{i=1}^{n-1} n - i.$$

Now, if we don’t like to deal with summing the values of $(n - i)$, we can observe that the values we are summing are $n - 1, n - 2, \dots, 1$, so we may write that

$$\sum_{i=1}^{n-1} n - i = \sum_{i=1}^{n-1} i.$$

A clever trick, usually attributed to Gauss, gives us a shorter formula for this sum.

We write

$$\begin{array}{cccccccc} 1 & + & 2 & + & \cdots & + & n - 2 & + & n - 1 \\ + & n - 1 & + & n - 2 & + & \cdots & + & 2 & + & 1 \\ \hline n & + & n & + & \cdots & + & n & + & n \end{array}$$

The sum below the horizontal line has $n - 1$ terms each equal to n , and thus it is $n(n - 1)$. It is the sum of the two sums above the line, and since these sums are equal (being identical except for being in reverse order), the sum below the line must be twice either sum above, so either of the sums above must be $n(n - 1)/2$. In other words, we may write

$$\sum_{i=1}^{n-1} n - i = \sum_{i=1}^{n-1} i = \frac{n(n - 1)}{2}.$$

This lovely trick gives us little or no real mathematical skill; learning how to think about things to discover answers ourselves is much more useful. After we analyze Exercise 1.1-2 and abstract the process we are using there, we will be able to come back to this problem at the end of this section and see a way that we could have discovered this formula for ourselves without any tricks.

The Product Principle

Exercise 1.1-2 The loop below is part of a program which computes the product of two matrices. (You don’t need to know what the product of two matrices is to answer this question.)

```

(1) for  $i = 1$  to  $r$ 
(2)     for  $j = 1$  to  $m$ 
(3)          $S = 0$ 
(4)         for  $k = 1$  to  $n$ 
(5)              $S = S + A[i, k] * B[k, j]$ 
(6)          $C[i, j] = S$ 

```

How many multiplications (expressed in terms of r , m , and n) does this code carry out in line 5?

Exercise 1.1-3 Consider the following longer piece of pseudocode that sorts a list of numbers and then counts “big gaps” in the list (for this problem, a big gap in the list is a place where a number in the list is more than twice the previous number):

```

(1) for  $i = 1$  to  $n - 1$ 
(2)      $minval = A[i]$ 
(3)      $minindex = i$ 
(4)     for  $j = i$  to  $n$ 
(5)         if ( $A[j] < minval$ )
(6)              $minval = A[j]$ 
(7)              $minindex = j$ 
(8)     exchange  $A[i]$  and  $A[minindex]$ 
(9)
(10) for  $i = 2$  to  $n$ 
(11)     if ( $A[i] > 2 * A[i - 1]$ )
(12)          $bigjump = bigjump + 1$ 

```

How many comparisons does the above code make in lines 5 and 11 ?

In Exercise 1.1-2, the program segment in lines 4 through 5, which we call the “inner loop,” takes exactly n steps, and thus makes n multiplications, regardless of what the variables i and j are. The program segment in lines 2 through 5 repeats the inner loop exactly m times, regardless of what i is. Thus this program segment makes n multiplications m times, so it makes nm multiplications.

Why did we add in Exercise 1.1-1, but multiply here? We can answer this question using the abstract point of view we adopted in discussing Exercise 1.1-1. Our algorithm performs a certain set of multiplications. For any given i , the set of multiplications performed in lines 2 through 5 can be divided into the set S_1 of multiplications performed when $j = 1$, the set S_2 of multiplications performed when $j = 2$, and, in general, the set S_j of multiplications performed for any given j value. Each set S_j consists of those multiplications the inner loop carries out for a particular value of j , and there are exactly n multiplications in this set. Let T_i be the set of multiplications that our program segment carries out for a certain i value. The set T_i is the union of the sets S_j ; restating this as an equation, we get

$$T_i = \bigcup_{j=1}^m S_j.$$

Then, by the sum principle, the size of the set T_i is the sum of the sizes of the sets S_j , and a sum of m numbers, each equal to n , is mn . Stated as an equation,

$$|T_i| = \left| \bigcup_{j=1}^m S_j \right| = \sum_{j=1}^m |S_j| = \sum_{j=1}^m n = mn. \quad (1.3)$$

Thus we are multiplying because multiplication is repeated addition!

From our solution we can extract a second principle that simply shortcuts the use of the sum principle.

Principle 1.3 (Product Principle) *The size of a union of m disjoint sets, each of size n , is mn .*

We now complete our discussion of Exercise 1.1-2. Lines 2 through 5 are executed once for each value of i from 1 to r . Each time those lines are executed, they are executed with a different i value, so the set of multiplications in one execution is disjoint from the set of multiplications in any other execution. Thus the set of all multiplications our program carries out is a union of r disjoint sets T_i of mn multiplications each. Then by the product principle, the set of all multiplications has size rmn , so our program carries out rmn multiplications.

Exercise 1.1-3 demonstrates that thinking about whether the sum or product principle is appropriate for a problem can help to decompose the problem into easily-solvable pieces. If you can decompose the problem into smaller pieces and solve the smaller pieces, then you either add or multiply solutions to solve the larger problem. In this exercise, it is clear that the number of comparisons in the program fragment is the sum of the number of comparisons in the first loop in lines 1 through 8 with the number of comparisons in the second loop in lines 10 through 12 (what two disjoint sets are we talking about here?). Further, the first loop makes $n(n+1)/2 - 1$ comparisons², and that the second loop has $n - 1$ comparisons, so the fragment makes $n(n+1)/2 - 1 + n - 1 = n(n+1)/2 + n - 2$ comparisons.

Two element subsets

Often, there are several ways to solve a problem. We originally solved Exercise 1.1-1 by using the sum principal, but it is also possible to solve it using the product principal. Solving a problem two ways not only increases our confidence that we have found the correct solution, but it also allows us to make new connections and can yield valuable insight.

Consider the set of comparisons made by the entire execution of the code in this exercise. When $i = 1$, j takes on every value from 2 to n . When $i = 2$, j takes on every value from 3 to n . Thus, for each two numbers i and j , we compare $A[i]$ and $A[j]$ exactly once in our loop. (The order in which we compare them depends on whether i or j is smaller.) Thus the number of comparisons we make is the same as the number of two element subsets of the set $\{1, 2, \dots, n\}$ ³. In how many ways can we choose two elements from this set? If we choose a first and second element, there are n ways to choose a first element, and for each choice of the first element, there are $n - 1$ ways to choose a second element. Thus the set of all such choices is the union of n sets

²To see why this is true, ask yourself first where the $n(n+1)/2$ comes from, and then why we subtracted one.

³The relationship between the set of comparisons and the set of two-element subsets of $\{1, 2, \dots, n\}$ is an example of a bijection, an idea which will be examined more in Section 1.2.

of size $n - 1$, one set for each first element. Thus it might appear that, by the product principle, there are $n(n - 1)$ ways to choose two elements from our set. However, what we have chosen is an *ordered pair*, namely a pair of elements in which one comes first and the other comes second. For example, we could choose 2 first and 5 second to get the ordered pair $(2, 5)$, or we could choose 5 first and 2 second to get the ordered pair $(5, 2)$. Since each pair of distinct elements of $\{1, 2, \dots, n\}$ can be ordered in two ways, we get twice as many ordered pairs as two element sets. Thus, since the number of ordered pairs is $n(n - 1)$, the number of two element subsets of $\{1, 2, \dots, n\}$ is $n(n - 1)/2$. Therefore the answer to Exercise 1.1-1 is $n(n - 1)/2$. This number comes up so often that it has its own name and notation. We call this number “ n choose 2” and denote it by $\binom{n}{2}$. To summarize, $\binom{n}{2}$ stands for the number of two element subsets of an n element set and equals $n(n - 1)/2$. Since one answer to Exercise 1.1-1 is $1 + 2 + \dots + n - 1$ and a second answer to Exercise 1.1-1 is $\binom{n}{2}$, this shows that

$$1 + 2 + \dots + n - 1 = \binom{n}{2} = \frac{n(n - 1)}{2}.$$

Important Concepts, Formulas, and Theorems

1. *Set.* A *set* is a collection of objects. In a set order is not important. Thus the set $\{A, B, C\}$ is the same as the set $\{A, C, B\}$. An element either is or is not in a set; it cannot be in a set more than once, even if we have a description of a set which names that element more than once.
2. *Disjoint.* Two sets are called *disjoint* when they have no elements in common.
3. *Mutually disjoint sets.* A set of sets $\{S_1, \dots, S_n\}$ is a family of *mutually disjoint sets*, if each two of the sets S_i are disjoint.
4. *Size of a set.* Given a set S , the size of S , denoted $|S|$, is the number of distinct elements in S .
5. *Sum Principle.* The size of a union of a family of mutually disjoint sets is the sum of the sizes of the sets. In other words, if S_1, S_2, \dots, S_n are disjoint sets, then

$$|S_1 \cup S_2 \cup \dots \cup S_n| = |S_1| + |S_2| + \dots + |S_n|.$$

To write this without the “dots” that indicate left-out material, we write

$$\left| \bigcup_{i=1}^n S_i \right| = \sum_{i=1}^n |S_i|.$$

6. *Partition of a set.* A partition of a set S is a set of mutually disjoint subsets (sometimes called blocks) of S whose union is S .
7. *Sum of first $n - 1$ numbers.*

$$\sum_{i=1}^n n - i = \sum_{i=1}^{n-1} i = \frac{n(n - 1)}{2}.$$

8. *Product Principle.* The size of a union of m disjoint sets, each of size n , is mn .
9. *Two element subsets.* $\binom{n}{2}$ stands for the number of two element subsets of an n element set and equals $n(n - 1)/2$. $\binom{n}{2}$ is read as “ n choose 2.”

Problems

1. The segment of code below is part of a program that uses insertion sort to sort a list A

```

for  $i = 2$  to  $n$ 
     $j = i$ 
    while  $j \geq 2$  and  $A[j] < A[j - 1]$ 
        exchange  $A[j]$  and  $A[j - 1]$ 
         $j --$ 

```

What is the maximum number of times (considering all lists of n items you could be asked to sort) the program makes the comparison $A[j] < A[j - 1]$? Describe as succinctly as you can those lists that require this number of comparisons.

2. Five schools are going to send their baseball teams to a tournament, in which each team must play each other team exactly once. How many games are required?
3. Use notation similar to that in Equations 1.2 and 1.3 to rewrite the solution to Exercise 1.1-3 more algebraically.
4. In how many ways can you draw a first card and then a second card from a deck of 52 cards?
5. In how many ways can you draw two cards from a deck of 52 cards.
6. In how many ways may you draw a first, second, and third card from a deck of 52 cards?
7. In how many ways may a ten person club select a president and a secretary-treasurer from among its members?
8. In how many ways may a ten person club select a two person executive committee from among its members?
9. In how many ways may a ten person club select a president and a two person executive advisory board from among its members (assuming that the president is not on the advisory board)?
10. By using the formula for $\binom{n}{2}$ it is straightforward to show that

$$n \binom{n-1}{2} = \binom{n}{2} (n-2).$$

However this proof just uses blind substitution and simplification. Find a more conceptual explanation of why this formula is true.

11. If M is an m element set and N is an n -element set, how many ordered pairs are there whose first member is in M and whose second member is in N ?
12. In the local ice cream shop, there are 10 different flavors. How many different two-scoop cones are there? (Following your mother's rule that it all goes to the same stomach, a cone with a vanilla scoop on top of a chocolate scoop is considered the same as a cone with a chocolate scoop on top of a vanilla scoop.)

13. Now suppose that you decide to disagree with your mother in Exercise 12 and say that the order of the scoops does matter. How many different possible two-scoop cones are there?
14. Suppose that on day 1 you receive 1 penny, and, for $i > 1$, on day i you receive twice as many pennies as you did on day $i - 1$. How many pennies will you have on day 20? How many will you have on day n ? Did you use the sum or product principal?
15. The “Pile High Deli” offers a “simple sandwich” consisting of your choice of one of five different kinds of bread with your choice of butter or mayonnaise or no spread, one of three different kinds of meat, and one of three different kinds of cheese, with the meat and cheese “piled high” on the bread. In how many ways may you choose a simple sandwich?
16. Do you see any unnecessary steps in the pseudocode of Exercise 1.1-3?

1.2 Counting Lists, Permutations, and Subsets.

Using the Sum and Product Principles

Exercise 1.2-1 A password for a certain computer system is supposed to be between 4 and 8 characters long and composed of lower and/or upper case letters. How many passwords are possible? What counting principles did you use? Estimate the percentage of the possible passwords that have exactly four characters.

A good way to attack a counting problem is to ask if we could use either the sum principle or the product principle to simplify or completely solve it. Here that question might lead us to think about the fact that a password can have 4, 5, 6, 7 or 8 characters. The set of all passwords is the union of those with 4, 5, 6, 7, and 8 letters so the sum principle might help us. To write the problem algebraically, let P_i be the set of i -letter passwords and P be the set of all possible passwords. Clearly,

$$P = P_4 \cup P_5 \cup P_6 \cup P_7 \cup P_8 .$$

The P_i are mutually disjoint, and thus we can apply the sum principle to obtain

$$|P| = \sum_{i=4}^8 |P_i| .$$

We still need to compute $|P_i|$. For an i -letter password, there are 52 choices for the first letter, 52 choices for the second and so on. Thus by the product principle, $|P_i|$, the number of passwords with i letters is 52^i . Therefore the total number of passwords is

$$52^4 + 52^5 + 52^6 + 52^7 + 52^8 .$$

Of these, 52^4 have four letters, so the percentage with 54 letters is

$$\frac{100 \cdot 52^4}{52^4 + 52^5 + 52^6 + 52^7 + 52^8} .$$

Although this is a nasty formula to evaluate by hand, we can get a quite good estimate as follows. Notice that 52^8 is 52 times as big as 52^7 , and even more dramatically larger than any other term in the sum in the denominator. Thus the ratio is just a bit less than

$$\frac{100 \cdot 52^4}{52^8} ,$$

which is $100/52^4$, or approximately .000014. Thus to five decimal places, only .00001% of the passwords have four letters. It is therefore much easier to guess a password that we know has four letters than it is to guess one that has between 4 and 8 letters—roughly 7 million times easier!

In our solution to Exercise 1.2-1, we casually referred to the use of the product principle in computing the number of passwords with i letters. We didn't write any set as a union of sets of equal size. We could have, but it would have been clumsy and repetitive. For this reason we will state a second version of the product principle that we can derive from the version for unions of sets by using the idea of mathematical induction that we study in Chapter 4.

Version 2 of the *product principle* states:

Principle 1.4 (Product Principle, Version 2) *If a set S of lists of length m has the properties that*

1. *There are i_1 different first elements of lists in S , and*
2. *For each $j > 1$ and each choice of the first $j - 1$ elements of a list in S there are i_j choices of elements in position j of those lists,*

then there are $i_1 i_2 \cdots i_m = \prod_{k=1}^m i_k$ lists in S .

Let's apply this version of the product principle to compute the number of m -letter passwords. Since an m -letter password is just a list of m letters, and since there are 52 different first elements of the password and 52 choices for each other position of the password, we have that $i_1 = 52$, $i_2 = 52, \dots, i_m = 52$. Thus, this version of the product principle tells us immediately that the number of passwords of length m is $i_1 i_2 \cdots i_m = 52^m$.

In our statement of version 2 of the Product Principle, we have introduced a new notation, the use of Π to stand for product. This notation is called the *product notation*, and it is used just like summation notation. In particular, $\prod_{k=1}^m i_k$ is read as "The product from $k = 1$ to m of i_k ." Thus $\prod_{k=1}^m i_k$ means the same thing as $i_1 \cdot i_2 \cdots i_m$.

Lists and functions

We have left a term undefined in our discussion of version 2 of the product principle, namely the word "list." A *list* of 3 things chosen from a set T consists of a first member t_1 of T , a second member t_2 of T , and a third member t_3 of T . If we rewrite the list in a different order, we get a different list. A list of k things chosen from T consists of a first member of T through a k th member of T . We can use the word "function," which you probably recall from algebra or calculus, to be more precise.

Recall that a function from a set S (called the *domain* of the function) to a set T (called the *range* of the function) is a relationship between the elements of S and the elements of T that relates exactly one element of T to each element of S . We use a letter like f to stand for a function and use $f(x)$ to stand for the one and only one element of T that the function relates to the element x of S . You are probably used to thinking of functions in terms of formulas like $f(x) = x^2$. We need to use formulas like this in algebra and calculus because the functions that you study in algebra and calculus have infinite sets of numbers as their domains and ranges. In discrete mathematics, functions often have finite sets as their domains and ranges, and so it is possible to describe a function by saying exactly what it is. For example

$$f(1) = \text{Sam}, f(2) = \text{Mary}, f(3) = \text{Sarah}$$

is a function that describes a list of three people. This suggests a precise definition of a list of k elements from a set T : A *list of k elements* from a set T is a function from $\{1, 2, \dots, k\}$ to T .

Exercise 1.2-2 Write down all the functions from the two-element set $\{1, 2\}$ to the two-element set $\{a, b\}$.

Exercise 1.2-3 How many functions are there from a two-element set to a three element set?

Exercise 1.2-4 How many functions are there from a three-element set to a two-element set?

In Exercise 1.2-2 one thing that is difficult is to choose a notation for writing the functions down. We will use f_1, f_2 , etc., to stand for the various functions we find. To describe a function f_i from $\{1, 2\}$ to $\{a, b\}$ we have to specify $f_i(1)$ and $f_i(2)$. We can write

$$\begin{aligned} f_1(1) &= a & f_1(2) &= b \\ f_2(1) &= b & f_2(2) &= a \\ f_3(1) &= a & f_3(2) &= a \\ f_4(1) &= b & f_4(2) &= b \end{aligned}$$

We have simply written down the functions as they occurred to us. How do we know we have all of them? The set of all functions from $\{1, 2\}$ to $\{a, b\}$ is the union of the functions f_i that have $f_i(1) = a$ and those that have $f_i(1) = b$. The set of functions with $f_i(1) = a$ has two elements, one for each choice of $f_i(2)$. Therefore by the product principle the set of all functions from $\{1, 2\}$ to $\{a, b\}$ has size $2 \cdot 2 = 4$.

To compute the number of functions from a two element set (say $\{1, 2\}$) to a three element set, we can again think of using f_i to stand for a typical function. Then the set of all functions is the union of three sets, one for each choice of $f_i(1)$. Each of these sets has three elements, one for each choice of $f_i(2)$. Thus by the product principle we have $3 \cdot 3 = 9$ functions from a two element set to a three element set.

To compute the number of functions from a three element set (say $\{1, 2, 3\}$) to a two element set, we observe that the set of functions is a union of four sets, one for each choice of $f_i(1)$ and $f_i(2)$ (as we saw in our solution to Exercise 1.2-2). But each of these sets has two functions in it, one for each choice of $f_i(3)$. Then by the product principle, we have $4 \cdot 2 = 8$ functions from a three element set to a two element set.

A function f is called *one-to-one* or an *injection* if whenever $x \neq y$, $f(x) \neq f(y)$. Notice that the two functions f_1 and f_2 we gave in our solution of Exercise 1.2-2 are one-to-one, but f_3 and f_4 are not.

A function f is called *onto* or a *surjection* if every element y in the range is $f(x)$ for some x in the domain. Notice that the functions f_1 and f_2 in our solution of Exercise 1.2-2 are onto functions but f_3 and f_4 are not.

Exercise 1.2-5 Using two-element sets or three-element sets as domains and ranges, find an example of a one-to-one function that is not onto.

Exercise 1.2-6 Using two-element sets or three-element sets as domains and ranges, find an example of an onto function that is not one-to-one.

Notice that the function given by $f(1) = c, f(2) = a$ is an example of a function from $\{1, 2\}$ to $\{a, b, c\}$ that is one-to one but not onto.

Also, notice that the function given by $f(1) = a, f(2) = b, f(3) = a$ is an example of a function from $\{1, 2, 3\}$ to $\{a, b\}$ that is onto but not one to one.

The Bijection Principle

Exercise 1.2-7 The loop below is part of a program to determine the number of triangles formed by n points in the plane.

```
(1) trianglecount = 0
(2) for i = 1 to n
(3)     for j = i + 1 to n
(4)         for k = j + 1 to n
(5)             if points i, j, and k are not collinear
(6)                 trianglecount = trianglecount + 1
```

How many times does the above code check three points to see if they are collinear in line 5?

In Exercise 1.2-7, we have a loop embedded in a loop that is embedded in another loop. Because the second loop, starting in line 3, begins with $j = i + 1$ and j increase up to n , and because the third loop, starting in line 4, begins with $k = j + 1$ and increases up to n , our code examines each triple of values i, j, k with $i < j < k$ exactly once. For example, if n is 4, then the triples (i, j, k) used by the algorithm, in order, are $(1, 2, 3)$, $(1, 2, 4)$, $(1, 3, 4)$, and $(2, 3, 4)$. Thus one way in which we might have solved Exercise 1.2-7 would be to compute the number of such triples, which we will call *increasing triples*. As with the case of two-element subsets earlier, the number of such triples is the number of three-element subsets of an n -element set. This is the second time that we have proposed counting the elements of one set (in this case the set of increasing triples chosen from an n -element set) by saying that it is equal to the number of elements of some other set (in this case the set of three element subsets of an n -element set). When are we justified in making such an assertion that two sets have the same size? There is another fundamental principle that abstracts our concept of what it means for two sets to have the same size. Intuitively two sets have the same size if we can match up their elements in such a way that each element of one set corresponds to exactly one element of the other set. This description carries with it some of the same words that appeared in the definitions of functions, one-to-one, and onto. Thus it should be no surprise that one-to-one and onto functions are part of our abstract principle.

Principle 1.5 (Bijection Principle) *Two sets have the same size if and only if there is a one-to-one function from one set onto the other.*

Our principle is called the *bijection principle* because a one-to-one and onto function is called a *bijection*. Another name for a bijection is a *one-to-one correspondence*. A bijection from a set to itself is called a *permutation* of that set.

What is the bijection that is behind our assertion that the number of increasing triples equals the number of three-element subsets? We define the function f to be the one that takes the increasing triple (i, j, k) to the subset $\{i, j, k\}$. Since the three elements of an increasing triple are different, the subset is a three element set, so we have a function from increasing triples to three element sets. Two different triples can't be the same set in two different orders, so different triples have to be associated with different sets. Thus f is one-to-one. Each set of three integers can be listed in increasing order, so it is the image under f of an increasing triple. Therefore f is onto. Thus we have a one-to-one correspondence, or bijection, between the set of increasing triples and the set of three element sets.

k -element permutations of a set

Since counting increasing triples is equivalent to counting three-element subsets, we can count increasing triples by counting three-element subsets instead. We use a method similar to the one we used to compute the number of two-element subsets of a set. Recall that the first step was to compute the number of ordered pairs of distinct elements we could choose from the set $\{1, 2, \dots, n\}$. So we now ask in how many ways may we choose an ordered triple of distinct elements from $\{1, 2, \dots, n\}$, or more generally, in how many ways may we choose a list of k distinct elements from $\{1, 2, \dots, n\}$. A list of k -distinct elements chosen from a set N is called a k -element permutation of N .⁴

How many 3-element permutations of $\{1, 2, \dots, n\}$ can we make? Recall that a k -element permutation is a list of k distinct elements. There are n choices for the first number in the list. For each way of choosing the first element, there are $n - 1$ choices for the second. For each choice of the first two elements, there are $n - 2$ ways to choose a third (distinct) number, so by version 2 of the product principle, there are $n(n - 1)(n - 2)$ ways to choose the list of numbers. For example, if n is 4, the three-element permutations of $\{1, 2, 3, 4\}$ are

$$L = \{123, 124, 132, 134, 142, 143, 213, 214, 231, 234, 241, 243, \\ 312, 314, 321, 324, 341, 342, 412, 413, 421, 423, 431, 432\}. \quad (1.4)$$

There are indeed $4 \cdot 3 \cdot 2 = 24$ lists in this set. Notice that we have listed the lists in the order that they would appear in a dictionary (assuming we treated numbers as we treat letters). This ordering of lists is called the *lexicographic ordering*.

A general pattern is emerging. To compute the number of k -element permutations of the set $\{1, 2, \dots, n\}$, we recall that they are lists and note that we have n choices for the first element of the list, and regardless of which choice we make, we have $n - 1$ choices for the second element of the list, and more generally, given the first $i - 1$ elements of a list we have $n - (i - 1) = n - i + 1$ choices for the i th element of the list. Thus by version 2 of the product principle, we have $n(n - 1) \cdots (n - k + 1)$ (which is the first k terms of $n!$) ways to choose a k -element permutation of $\{1, 2, \dots, n\}$. There is a very handy notation for this product first suggested by Don Knuth. We use $n^{\underline{k}}$ to stand for $n(n - 1) \cdots (n - k + 1) = \prod_{i=0}^{k-1} n - i$, and call it the k th falling factorial power of n . We can summarize our observations in a theorem.

Theorem 1.1 *The number k -element permutations of an n -element set is*

$$n^{\underline{k}} = \prod_{i=0}^{k-1} n - i = n(n - 1) \cdots (n - k + 1) = n! / (n - k)! .$$

Counting subsets of a set

We now return to the question of counting the number of three element subsets of a $\{1, 2, \dots, n\}$. We use $\binom{n}{3}$, which we read as “ n choose 3” to stand for the number of three element subsets of

⁴In particular a k -element permutation of $\{1, 2, \dots, k\}$ is a list of k distinct elements of $\{1, 2, \dots, k\}$, which, by our definition of a list is a function from $\{1, 2, \dots, k\}$ to $\{1, 2, \dots, k\}$. This function must be one-to-one since the elements of the list are distinct. Since there are k distinct elements of the list, every element of $\{1, 2, \dots, k\}$ appears in the list, so the function is onto. Therefore it is a bijection. Thus our definition of a permutation of a set is consistent with our definition of a k -element permutation in the case where the set is $\{1, 2, \dots, k\}$.

$\{1, 2, \dots, n\}$, or more generally of any n -element set. We have just carried out the first step of computing $\binom{n}{3}$ by counting the number of three-element permutations of $\{1, 2, \dots, n\}$.

Exercise 1.2-8 Let L be the set of all three-element permutations of $\{1, 2, 3, 4\}$, as in Equation 1.4. How many of the lists (permutations) in L are lists of the 3 element set $\{1, 3, 4\}$? What are these lists?

We see that this set appears in L as 6 different lists: 134, 143, 314, 341, 413, and 431. In general given three different numbers with which to create a list, there are three ways to choose the first number in the list, given the first there are two ways to choose the second, and given the first two there is only one way to choose the third element of the list. Thus by version 2 of the product principle once again, there are $3 \cdot 2 \cdot 1 = 6$ ways to make the list.

Since there are $n(n-1)(n-2)$ permutations of an n -element set, and each three-element subset appears in exactly 6 of these lists, the number of three-element permutations is six times the number of three element subsets. That is, $n(n-1)(n-2) = \binom{n}{3} \cdot 6$. Whenever we see that one number that counts something is the product of two other numbers that count something, we should expect that there is an argument using the product principle that explains why. Thus we should be able to see how to break the set of all 3-element permutations of $\{1, 2, \dots, n\}$ into either 6 disjoint sets of size $\binom{n}{3}$ or into $\binom{n}{3}$ subsets of size six. Since we argued that each three element subset corresponds to six lists, we have described how to get a set of six lists from one three-element set. Two different subsets could never give us the same lists, so our sets of three-element lists are disjoint. In other words, we have divided the set of all three-element permutations into $\binom{n}{3}$ mutually sets of size six. In this way the product principle does explain why $n(n-1)(n-2) = \binom{n}{3} \cdot 6$. By division we get that we have

$$\binom{n}{3} = n(n-1)(n-2)/6$$

three-element subsets of $\{1, 2, \dots, n\}$. For $n = 4$, the number is $4(3)(2)/6 = 4$. These sets are $\{1, 2, 3\}$, $\{1, 2, 4\}$, $\{1, 3, 4\}$, and $\{2, 3, 4\}$. It is straightforward to verify that each of these sets appears 6 times in L , as 6 different lists.

Essentially the same argument gives us the number of k -element subsets of $\{1, 2, \dots, n\}$. We denote this number by $\binom{n}{k}$, and read it as “ n choose k .” Here is the argument: the set of all k -element permutations of $\{1, 2, \dots, n\}$ can be partitioned into $\binom{n}{k}$ disjoint blocks⁵, each block consisting of all k -element permutations of a k -element subset of $\{1, 2, \dots, n\}$. But the number of k -element permutations of a k -element set is $k!$, either by version 2 of the product principle or by Theorem 1.1. Thus by version 1 of the product principle we get the equation

$$n^k = \binom{n}{k} k!.$$

Division by $k!$ gives us our next theorem.

Theorem 1.2 For integers n and k with $0 \leq k \leq n$, the number of k element subsets of an n element set is

$$\frac{n^k}{k!} = \frac{n!}{k!(n-k)!}$$

⁵Here we are using the language introduced for partitions of sets in Section 1.1

Proof: The proof is given above, except in the case that k is 0; however the only subset of our n -element set of size zero is the empty set, so we have exactly one such subset. This is exactly what the formula gives us as well. (Note that the cases $k = 0$ and $k = n$ both use the fact that $0! = 1$.⁶) The equality in the theorem comes from the definition of n^k . ■

Another notation for the numbers $\binom{n}{k}$ is $C(n, k)$. Thus we have that

$$C(n, k) = \binom{n}{k} = \frac{n!}{k!(n-k)!}. \quad (1.5)$$

These numbers are called *binomial coefficients* for reasons that will become clear later.

Important Concepts, Formulas, and Theorems

1. *List.* A list of k items chosen from a set X is a function from $\{1, 2, \dots, k\}$ to X .
2. *Lists versus sets.* In a list, the order in which elements appear in the list matters, and an element may appear more than once. In a set, the order in which we write down the elements of the set does not matter, and an element can appear at most once.
3. *Product Principle, Version 2.* If a set S of lists of length m has the properties that
 - (a) There are i_1 different first elements of lists in S , and
 - (b) For each $j > 1$ and each choice of the first $j - 1$ elements of a list in S there are i_j choices of elements in position j of those lists,

then there are $i_1 i_2 \cdots i_m$ lists in S .

4. *Product Notation.* We use the Greek letter Π to stand for product just as we use the Greek letter Σ to stand for sum. This notation is called the *product notation*, and it is used just like summation notation. In particular, $\prod_{k=1}^m i_k$ is read as “The product from $k = 1$ to m of i_k .” Thus $\prod_{k=1}^m i_k$ means the same thing as $i_1 \cdot i_2 \cdots i_m$.
5. *Function.* A *function* f from a set S to a set T is a relationship between S and T that relates exactly one element of T to each element of S . We write $f(x)$ for the one and only one element of T that the function f relates to the element x of S . The same element of T may be related to different members of S .
6. *Onto, Surjection* A function f from a set S to a set T is *onto* if for each element $y \in T$, there is at least one $x \in S$ such that $f(x) = y$. An onto function is also called a *surjection*.
7. *One-to-one, Injection.* A function f from a set S to a set T is *one-to-one* if, for each $x \in S$ and $y \in S$ with $x \neq y$, $f(x) \neq f(y)$. A one-to-one function is also called an *injection*.
8. *Bijection, One-to-one correspondence.* A function from a set S to a set T is a *bijection* if it is both one-to-one and onto. A bijection is sometimes called a *one-to-one correspondence*.
9. *Permutation.* A one-to-one function from a set S to S is called a *permutation* of S .

⁶There are many reasons why $0!$ is defined to be one; making the formula for $\binom{n}{k}$ work out is one of them.

10. *k*-element permutation. A *k*-element permutation of a set S is a list of k distinct elements of S .
11. *k*-element subsets. *n* choose *k*. *Binomial Coefficients*. For integers n and k with $0 \leq k \leq n$, the number of k element subsets of an n element set is $n!/k!(n-k)!$. The number of k -element subsets of an n -element set is usually denoted by $\binom{n}{k}$ or $C(n, k)$, both of which are read as “ n choose k .” These numbers are called *binomial coefficients*.
12. The number of k -element permutations of an n -element set is

$$n^{\underline{k}} = n(n-1) \cdots (n-k+1) = n!/(n-k)!$$

13. When we have a formula to count something and the formula expresses the result as a product, it is useful to try to understand whether and how we could use the product principle to prove the formula.

Problems

1. The “Pile High Deli” offers a “simple sandwich” consisting of your choice of one of five different kinds of bread with your choice of butter or mayonnaise or no spread, one of three different kinds of meat, and one of three different kinds of cheese, with the meat and cheese “piled high” on the bread. In how many ways may you choose a simple sandwich?
2. In how many ways can we pass out k distinct pieces of fruit to n children (with no restriction on how many pieces of fruit a child may get)?
3. Write down all the functions from the three-element set $\{1, 2, 3\}$ to the set $\{a, b\}$. Indicate which functions, if any, are one-to-one. Indicate which functions, if any, are onto.
4. Write down all the functions from the two element set $\{1, 2\}$ to the three element set $\{a, b, c\}$. Indicate which functions, if any, are one-to-one. Indicate which functions, if any, are onto.
5. There are more functions from the real numbers to the real numbers than most of us can imagine. However in discrete mathematics we often work with functions from a finite set S with s elements to a finite set T with t elements. Then there are only a finite number of functions from S to T . How many functions are there from S to T in this case?
6. Assuming $k \leq n$, in how many ways can we pass out k distinct pieces of fruit to n children if each child may get at most one? What is the number if $k > n$? Assume for both questions that we pass out all the fruit.
7. Assume $k \leq n$, in how many ways can we pass out k identical pieces of fruit to n children if each child may get at most one? What is the number if $k > n$? Assume for both questions that we pass out all the fruit.
8. What is the number of five digit (base ten) numbers? What is the number of five digit numbers that have no two consecutive digits equal? What is the number that have at least one pair of consecutive digits equal?

9. We are making a list of participants in a panel discussion on allowing alcohol on campus. They will be sitting behind a table in the order in which we list them. There will be four administrators and four students. In how many ways may we list them if the administrators must sit together in a group and the students must sit together in a group? In how many ways may we list them if we must alternate students and administrators?
10. (This problem is for students who are working on the relationship between k -element permutations and k -element subsets.) Write down all three element permutations of the five element set $\{1, 2, 3, 4, 5\}$ in lexicographic order. Underline those that correspond to the set $\{1, 3, 5\}$. Draw a rectangle around those that correspond to the set $\{2, 4, 5\}$. How many three-element permutations of $\{1, 2, 3, 4, 5\}$ correspond to a given 3-element set? How many three-element subsets does the set $\{1, 2, 3, 4, 5\}$ have?
11. In how many ways may a class of twenty students choose a group of three students from among themselves to go to the professor and explain that the three-hour labs are actually taking ten hours?
12. We are choosing participants for a panel discussion allowing on allowing alcohol on campus. We have to choose four administrators from a group of ten administrators and four students from a group of twenty students. In how many ways may we do this?
13. We are making a list of participants in a panel discussion on allowing alcohol on campus. They will be sitting behind a table in the order in which we list them. There will be four administrators chosen from a group of ten administrators and four students chosen from a group of twenty students. In how many ways may we choose and list them if the administrators must sit together in a group and the students must sit together in a group? In how many ways may we choose and list them if we must alternate students and administrators?
14. In the local ice cream shop, you may get a sundae with two scoops of ice cream from 10 flavors (in accordance with your mother's rules from Problem 12 in Section 1.1, the way the scoops sit in the dish does not matter), any one of three flavors of topping, and any (or all or none) of whipped cream, nuts and a cherry. How many different sundaes are possible?
15. In the local ice cream shop, you may get a three-way sundae with three of the ten flavors of ice cream, any one of three flavors of topping, and any (or all or none) of whipped cream, nuts and a cherry. How many different sundaes are possible(in accordance with your mother's rules from Problem 12 in Section 1.1, the way the scoops sit in the dish does not matter) ?
16. A tennis club has $2n$ members. We want to pair up the members by twos for singles matches. In how many ways may we pair up all the members of the club? Suppose that in addition to specifying who plays whom, for each pairing we say who serves first. Now in how many ways may we specify our pairs?
17. A basketball team has 12 players. However, only five players play at any given time during a game. In how many ways may the coach choose the five players? To be more realistic, the five players playing a game normally consist of two guards, two forwards, and one center. If there are five guards, four forwards, and three centers on the team, in how many ways can the coach choose two guards, two forwards, and one center? What if one of the centers is equally skilled at playing forward?

18. Explain why a function from an n -element set to an n -element set is one-to-one if and only if it is onto.
19. The function g is called an *inverse* to the function f if the domain of g is the range of f , if $g(f(x)) = x$ for every x in the domain of f and if $f(g(y)) = y$ for each y in the range of f .
 - (a) Explain why a function is a bijection if and only if it has an inverse function.
 - (b) Explain why a function that has an inverse function has only one inverse function.

1.3 Binomial Coefficients

In this section, we will explore various properties of binomial coefficients. Remember that we defined the quantity $\binom{n}{k}$ to be the number of k -element subsets of an n -element set.

Pascal's Triangle

Table 1 contains the values of the binomial coefficients $\binom{n}{k}$ for $n = 0$ to 6 and all relevant k values. The table begins with a 1 for $n = 0$ and $k = 0$, because the empty set, the set with no elements, has exactly one 0-element subset, namely itself. We have not put any value into the table for a value of k larger than n , because we haven't directly said what we mean by the binomial coefficient $\binom{n}{k}$ in that case. However, since there are no subsets of an n -element set that have size larger than n , it is natural to say that $\binom{n}{k}$ is zero when $k > n$. Therefore we define $\binom{n}{k}$ to be zero⁷ when $k > n$. Thus we could fill in the empty places in the table with zeros. The table is easier to read if we don't fill in the empty spaces, so we just remember that they are zero.

Table 1.1: A table of binomial coefficients

$n \setminus k$	0	1	2	3	4	5	6
0	1						
1	1	1					
2	1	2	1				
3	1	3	3	1			
4	1	4	6	4	1		
5	1	5	10	10	5	1	
6	1	6	15	20	15	6	1

Exercise 1.3-1 What general properties of binomial coefficients do you see in Table 1.1

Exercise 1.3-2 What is the next row of the table of binomial coefficients?

Several properties of binomial coefficients are apparent in Table 1.1. Each row begins with a 1, because $\binom{n}{0}$ is always 1. This is the case because there is just one subset of an n -element set with 0 elements, namely the empty set. Similarly, each row ends with a 1, because an n -element set S has just one n -element subset, namely S itself. Each row increases at first, and then decreases. Further the second half of each row is the reverse of the first half. The array of numbers called *Pascal's Triangle* emphasizes that symmetry by rearranging the rows of the table so that they line up at their centers. We show this array in Table 2. When we write down Pascal's triangle, we leave out the values of n and k .

You may know a method for creating Pascal's triangle that does not involve computing binomial coefficients, but rather creates each row from the row above. Each entry in Table 1.2, except for the ones, is the sum of the entry directly above it to the left and the entry directly

⁷If you are thinking "But we did define $\binom{n}{k}$ to be zero when $k > n$ by saying that it is the number of k element subsets of an n -element set, so of course it is zero," then good for you.

Table 1.2: Pascal's Triangle

				1			
				1		1	
			1	2		1	
		1	3	3		1	
	1	4	6	4		1	
	1	5	10	10		5	1
1	6	15	20	15		6	1

above it to the right. We call this the *Pascal Relationship*, and it gives another way to compute binomial coefficients without doing the multiplying and dividing in Equation 1.5. If we wish to compute many binomial coefficients, the Pascal relationship often yields a more efficient way to do so. Once the coefficients in a row have been computed, the coefficients in the next row can be computed using only one addition per entry.

We now verify that the two methods for computing Pascal's triangle always yield the same result. In order to do so, we need an algebraic statement of the Pascal Relationship. In Table 1.1, each entry is the sum of the one above it and the one above it and to the left. In algebraic terms, then, the Pascal Relationship says

$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}, \quad (1.6)$$

whenever $n > 0$ and $0 < k < n$. It is possible to give a purely algebraic (and rather dreary) proof of this formula by plugging in our earlier formula for binomial coefficients into all three terms and verifying that we get an equality. A guiding principle of discrete mathematics is that when we have a formula that relates the numbers of elements of several sets, we should find an explanation that involves a relationship among the sets.

A proof using the Sum Principle

From Theorem 1.2 and Equation 1.5, we know that the expression $\binom{n}{k}$ is the number of k -element subsets of an n element set. Each of the three terms in Equation 1.6 therefore represents the number of subsets of a particular size chosen from an appropriately sized set. In particular, the three terms are the number of k -element subsets of an n -element set, the number of $(k-1)$ -element subsets of an $(n-1)$ -element set, and the number of k -element subsets of an $(n-1)$ -element set. We should, therefore, be able to explain the relationship among these three quantities using the sum principle. This explanation will provide a proof, just as valid a proof as an algebraic derivation. Often, a proof using the sum principle will be less tedious, and will yield more insight into the problem at hand.

Before giving such a proof in Theorem 1.3 below, we work out a special case. Suppose $n = 5$, $k = 2$. Equation 1.6 says that

$$\binom{5}{2} = \binom{4}{1} + \binom{4}{2}. \quad (1.7)$$

Because the numbers are small, it is simple to verify this by using the formula for binomial coefficients, but let us instead consider subsets of a 5-element set. Equation 1.7 says that the number of 2 element subsets of a 5 element set is equal to the number of 1 element subsets of a 4 element set plus the number of 2 element subsets of a 4 element set. But to apply the sum principle, we would need to say something stronger. To apply the sum principle, we should be able to partition the set of 2 element subsets of a 5 element set into 2 disjoint sets, one of which has the same size as the number of 1 element subsets of a 4 element set and one of which has the same size as the number of 2 element subsets of a 4 element set. Such a partition provides a proof of Equation 1.7. Consider now the set $S = \{A, B, C, D, E\}$. The set of two element subsets is

$$S_1 = \{\{A, B\}, \{AC\}, \{A, D\}, \{A, E\}, \{B, C\}, \{B, D\}, \{B, E\}, \{C, D\}, \{C, E\}, \{D, E\}\}.$$

We now partition S_1 into 2 blocks, S_2 and S_3 . S_2 contains all sets in S_1 that do contain the element E , while S_3 contains all sets in S_1 that do not contain the element E . Thus,

$$S_2 = \{\{AE\}, \{BE\}, \{CE\}, \{DE\}\}$$

and

$$S_3 = \{\{AB\}, \{AC\}, \{AD\}, \{BC\}, \{BD\}, \{CD\}\}.$$

Each set in S_2 must contain E and thus contains one other element from S . Since there are 4 other elements in S that we can choose along with E , we have $|S_2| = \binom{4}{1}$. Each set in S_3 contains 2 elements from the set $\{A, B, C, D\}$. There are $\binom{4}{2}$ ways to choose such a two-element subset of $\{A < B < C < D\}$. But $S_1 = S_2 \cup S_3$ and S_2 and S_3 are disjoint, and so, by the sum principle, Equation 1.7 must hold.

We now give a proof for general n and k .

Theorem 1.3 *If n and k are integers with $n > 0$ and $0 < k < n$, then*

$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}.$$

Proof: The formula says that the number of k -element subsets of an n -element set is the sum of two numbers. As in our example, we will apply the sum principle. To apply it, we need to represent the set of k -element subsets of an n -element set as a union of two other disjoint sets. Suppose our n -element set is $S = \{x_1, x_2, \dots, x_n\}$. Then we wish to take S_1 , say, to be the $\binom{n}{k}$ -element set of all k -element subsets of S and partition it into two disjoint sets of k -element subsets, S_2 and S_3 , where the sizes of S_2 and S_3 are $\binom{n-1}{k-1}$ and $\binom{n-1}{k}$ respectively. We can do this as follows. Note that $\binom{n-1}{k}$ stands for the number of k element subsets of the first $n-1$ elements x_1, x_2, \dots, x_{n-1} of S . Thus we can let S_3 be the set of k -element subsets of S that don't contain x_n . Then the only possibility for S_2 is the set of k -element subsets of S that *do* contain x_n . How can we see that the number of elements of this set S_2 is $\binom{n-1}{k-1}$? By observing that removing x_n from each of the elements of S_2 gives a $(k-1)$ -element subset of $S' = \{x_1, x_2, \dots, x_{n-1}\}$. Further each $(k-1)$ -element subset of S' arises in this way from one and only one k -element subset of S containing x_n . Thus the number of elements of S_2 is the number of $(k-1)$ -element subsets

of S' , which is $\binom{n-1}{k-1}$. Since S_2 and S_3 are two disjoint sets whose union is S , the sum principle shows that the number of elements of S is $\binom{n-1}{k-1} + \binom{n-1}{k}$. ■

Notice that in our proof, we used a bijection that we did not explicitly describe. Namely, there is a bijection f between S_3 (the k -element sets of S that contain x_n) and the $(k-1)$ -element subsets of S' . For any subset K in S_3 , We let $f(K)$ be the set we obtain by removing x_n from K . It is immediate that this is a bijection, and so the bijection principle tells us that the size of S_3 is the size of the set of all subsets of S' .

The Binomial Theorem

Exercise 1.3-3 What is $(x+y)^3$? What is $(x+1)^4$? What is $(2+y)^4$? What is $(x+y)^4$?

The number of k -element subsets of an n -element set is called a *binomial coefficient* because of the role that these numbers play in the algebraic expansion of a binomial $x+y$. The **Binomial Theorem** states that

Theorem 1.4 (Binomial Theorem) For any integer $n \geq 0$

$$(x+y)^n = \binom{n}{0}x^n + \binom{n}{1}x^{n-1}y + \binom{n}{2}x^{n-2}y^2 + \cdots + \binom{n}{n-1}xy^{n-1} + \binom{n}{n}y^n, \quad (1.8)$$

or in summation notation,

$$(x+y)^n = \sum_{i=0}^n \binom{n}{i} x^{n-i} y^i.$$

Unfortunately when most people first see this theorem, they do not have the tools to see easily why it is true. Armed with our new methodology of using subsets to prove algebraic identities, we can give a proof of this theorem.

Let us begin by considering the example $(x+y)^3$ which by the binomial theorem is

$$(x+y)^3 = \binom{3}{0}x^3 + \binom{3}{1}x^2y + \binom{3}{2}xy^2 + \binom{3}{3}y^3 \quad (1.9)$$

$$= x^3 + 3x^2y + 3xy^2 + y^3. \quad (1.10)$$

Suppose that we did not know the binomial theorem but still wanted to compute $(x+y)^3$. Then we would write out $(x+y)(x+y)(x+y)$ and perform the multiplication. Probably we would multiply the first two terms, obtaining $x^2 + 2xy + y^2$, and then multiply this expression by $x+y$. Notice that by applying distributive laws you get

$$(x+y)(x+y) = (x+y)x + (x+y)y = xx + xy + yx + y. \quad (1.11)$$

We could use the commutative law to put this into the usual form, but let us hold off for a moment so we can see a pattern evolve. To compute $(x+y)^3$, we can multiply the expression on the right hand side of Equation 1.11 by $x+y$ using the distributive laws to get

$$(xx + xy + yx + yy)(x+y) = (xx + xy + yx + yy)x + (xx + xy + yx + yy)y \quad (1.12)$$

$$= xxx + xyx + yxx + yxx + xxy + xyy + yxy + yyy \quad (1.13)$$

Each of these 8 terms that we got from the distributive law may be thought of as a product of terms, one from the first binomial, one from the second binomial, and one from the third binomial. Multiplication is commutative, so many of these products are the same. In fact, we have one xxx or x^3 product, three products with two x 's and one y , or x^2y , three products with one x and two y 's, or xy^2 and one product which becomes y^3 . Now look at Equation 1.9, which summarizes this process. There are $\binom{3}{0} = 1$ way to choose a product with 3 x 's and 0 y 's, $\binom{3}{1} = 3$ way to choose a product with 2 x 's and 1 y , etc. Thus we can understand the binomial theorem as counting the subsets of our binomial factors from which we choose a y -term to get a product with k y 's in multiplying a string of n binomials.

Essentially the same explanation gives us a proof of the binomial theorem. Note that when we multiplied out three factors of $(x + y)$ using the distributive law but not collecting like terms, we had a sum of eight products. Each factor of $(x + y)$ doubles the number of summands. Thus when we apply the distributive law as many times as possible (without applying the commutative law and collecting like terms) to a product of n binomials all equal to $(x + y)$, we get 2^n summands. Each summand is a product of a length n list of x 's and y 's. In each list, the i th entry comes from the i th binomial factor. A list that becomes $x^{n-k}y^k$ when we use the commutative law will have a y in k of its places and an x in the remaining places. The number of lists that have a y in k places is thus the number of ways to select k binomial factors to contribute a y to our list. But the number of ways to select k binomial factors from n binomial factors is simply $\binom{n}{k}$, and so that is the coefficient of $x^{n-k}y^k$. This proves the binomial theorem.

Applying the Binomial Theorem to the remaining questions in Exercise 1.3-3 gives us

$$\begin{aligned}(x + 1)^4 &= x^4 + 4x^3 + 6x^2 + 4x + 1 \\ (2 + y)^4 &= 16 + 32y + 24y^2 + 8y^3 + y^4 \text{ and} \\ (x + y)^4 &= x^4 + 4x^3y + 6x^2y^2 + 4xy^3 + y^4.\end{aligned}$$

Labeling and trinomial coefficients

Exercise 1.3-4 Suppose that I have k labels of one kind and $n - k$ labels of another. In how many different ways may I apply these labels to n objects?

Exercise 1.3-5 Show that if we have k_1 labels of one kind, k_2 labels of a second kind, and $k_3 = n - k_1 - k_2$ labels of a third kind, then there are $\frac{n!}{k_1!k_2!k_3!}$ ways to apply these labels to n objects.

Exercise 1.3-6 What is the coefficient of $x^{k_1}y^{k_2}z^{k_3}$ in $(x + y + z)^n$?

Exercise 1.3-4 and Exercise 1.3-5 can be thought of as immediate applications of binomial coefficients. For Exercise 1.3-4, there are $\binom{n}{k}$ ways to choose the k objects that get the first label, and the other objects get the second label, so the answer is $\binom{n}{k}$. For Exercise 1.3-5, there are $\binom{n}{k_1}$ ways to choose the k_1 objects that get the first kind of label, and then there are $\binom{n-k_1}{k_2}$ ways to choose the objects that get the second kind of label. After that, the remaining $k_3 = n - k_1 - k_2$ objects get the third kind of label. The total number of labellings is thus, by the product principle, the product of the two binomial coefficients, which simplifies as follows.

$$\binom{n}{k_1} \binom{n-k_1}{k_2} = \frac{n!}{k_1!(n-k_1)!} \frac{(n-k_1)!}{k_2!(n-k_1-k_2)!}$$

$$\begin{aligned}
&= \frac{n!}{k_1!k_2!(n-k_1-k_2)!} \\
&= \frac{n!}{k_1!k_2!k_3!}.
\end{aligned}$$

A more elegant approach to Exercise 1.3-4, Exercise 1.3-5, and other related problems appears in the next section.

Exercise 1.3-6 shows how Exercise 1.3-5 applies to computing powers of trinomials. In expanding $(x + y + z)^n$, we think of writing down n copies of the trinomial $x + y + z$ side by side, and applying the distributive laws until we have a sum of terms each of which is a product of x 's, y 's and z 's. How many such terms do we have with k_1 x 's, k_2 y 's and k_3 z 's? Imagine choosing x from some number k_1 of the copies of the trinomial, choosing y from some number k_2 , and z from the remaining k_3 copies, multiplying all the chosen terms together, and adding up over all ways of picking the k_i s and making our choices. Choosing x from a copy of the trinomial “labels” that copy with x , and the same for y and z , so the number of choices that yield $x^{k_1}y^{k_2}z^{k_3}$ is the number of ways to label n objects with k_1 labels of one kind, k_2 labels of a second kind, and k_3 labels of a third. Notice that this requires that $k_3 = n - k_1 - k_2$. By analogy with our notation for a binomial coefficient, we define the *trinomial coefficient* $\binom{n}{k_1, k_2, k_3}$ to be $\frac{n!}{k_1!k_2!k_3!}$ if $k_1 + k_2 + k_3 = n$ and 0 otherwise. Then $\binom{n}{k_1, k_2, k_3}$ is the coefficient of $x^{k_1}y^{k_2}z^{k_3}$ in $(x + y + z)^n$. This is sometimes called the *trinomial theorem*.

Important Concepts, Formulas, and Theorems

1. *Pascal Relationship*. The Pascal Relationship says that

$$\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k},$$

whenever $n > 0$ and $0 < k < n$.

2. *Pascal's Triangle*. Pascal's Triangle is the triangular array of numbers we get by putting ones in row n and column 0 and in row n and column n of a table for every positive integer n and then filling the remainder of the table by letting the number in row n and column j be the sum of the numbers in row $n - 1$ and columns $j - 1$ and j whenever $0 < j < n$.

3. *Binomial Theorem*. The **Binomial Theorem** states that for any integer $n \geq 0$

$$(x + y)^n = x^n + \binom{n}{1}x^{n-1}y + \binom{n}{2}x^{n-2}y^2 + \cdots + \binom{n}{n-1}xy^{n-1} + \binom{n}{n}y^n,$$

or in summation notation,

$$(x + y)^n = \sum_{i=0}^n \binom{n}{i} x^{n-i} y^i.$$

4. *Labeling*. The number of ways to apply k labels of one kind and $n - k$ labels of another kind to n objects is $\binom{n}{k}$.
5. *Trinomial coefficient*. We define the *trinomial coefficient* $\binom{n}{k_1, k_2, k_3}$ to be $\frac{n!}{k_1!k_2!k_3!}$ if $k_1 + k_2 + k_3 = n$ and 0 otherwise.
6. *Trinomial Theorem*. The coefficient of $x^i y^j z^k$ in $(x + y + z)^n$ is $\binom{n}{i, j, k}$.

Problems

1. Find $\binom{12}{3}$ and $\binom{12}{9}$. What can you say in general about $\binom{n}{k}$ and $\binom{n}{n-k}$?
2. Find the row of the Pascal triangle that corresponds to $n = 8$.
3. Find the following
 - a. $(x + 1)^5$
 - b. $(x + y)^5$
 - c. $(x + 2)^5$
 - d. $(x - 1)^5$
4. Carefully explain the proof of the binomial theorem for $(x + y)^4$. That is, explain what each of the binomial coefficients in the theorem stands for and what powers of x and y are associated with them in this case.
5. If I have ten distinct chairs to paint, in how many ways may I paint three of them green, three of them blue, and four of them red? What does this have to do with labellings?
6. When n_1, n_2, \dots, n_k are nonnegative integers that add to n , the number $\frac{n!}{n_1!n_2!\dots n_k!}$ is called a *multinomial coefficient* and is denoted by $\binom{n}{n_1, n_2, \dots, n_k}$. A polynomial of the form $x_1 + x_2 + \dots + x_k$ is called a multinomial. Explain the relationship between powers of a multinomial and multinomial coefficients. This relationship is called the Multinomial Theorem.
7. Give a bijection that proves your statement about $\binom{n}{k}$ and $\binom{n}{n-k}$ in Problem 1 of this section.
8. In a Cartesian coordinate system, how many paths are there from the origin to the point with integer coordinates (m, n) if the paths are built up of exactly $m + n$ horizontal and vertical line segments each of length one?
9. What is the formula we get for the binomial theorem if, instead of analyzing the number of ways to choose k distinct y 's, we analyze the number of ways to choose k distinct x 's?
10. Explain the difference between choosing four disjoint three element sets from a twelve element set and labelling a twelve element set with three labels of type 1, three labels of type two, three labels of type 3, and three labels of type 4. What is the number of ways of choosing three disjoint four element subsets from a twelve element set? What is the number of ways of choosing four disjoint three element subsets from a twelve element set?
11. A 20 member club must have a President, Vice President, Secretary and Treasurer as well as a three person nominations committee. If the officers must be different people, and if no officer may be on the nominating committee, in how many ways could the officers and nominating committee be chosen? Answer the same question if officers may be on the nominating committee.
12. Prove Equation 1.6 by plugging in the formula for $\binom{n}{k}$.

13. Give two proofs that

$$\binom{n}{k} = \binom{n}{n-k}.$$

14. Give at least two proofs that

$$\binom{n}{k} \binom{k}{j} = \binom{n}{j} \binom{n-j}{k-j}.$$

15. Give at least two proofs that

$$\binom{n}{k} \binom{n-k}{j} = \binom{n}{j} \binom{n-j}{k}.$$

16. You need not compute all of rows 7, 8, and 9 of Pascal's triangle to use it to compute
- $\binom{9}{6}$
- . Figure out which entries of Pascal's triangle not given in Table 2 you actually need, and compute them to get
- $\binom{9}{6}$
- .

17. Explain why

$$\sum_{i=0}^n (-1)^i \binom{n}{i} = 0$$

18. Apply calculus and the binomial theorem to
- $(1+x)^n$
- to show that

$$\binom{n}{1} + 2\binom{n}{2} + 3\binom{n}{3} + \cdots = n2^{n-1}.$$

19. True or False:
- $\binom{n}{k} = \binom{n-2}{k-2} + \binom{n-2}{k-1} + \binom{n-2}{k}$
- . If true, give a proof. If false, give a value of
- n
- and
- k
- that show the statement is false, find an analogous true statement, and prove it.

1.4 Equivalence Relations and Counting (Optional)

The Symmetry Principle

Consider again the example from Section 1.2 in which we wanted to count the number of 3 element subsets of a four element set. To do so, we first formed all possible lists of $k = 3$ distinct elements chosen from an $n = 4$ element set. (See Equation 1.4.) The number of lists of k distinct elements is $n^k = n!/(n - k)!$. We then observed that two lists are equivalent as sets, if one can be obtained by rearranging (or “permuting”) the other. This process divides the lists up into classes, called *equivalence classes*, each of size $k!$. Returning to our example in Section 1.2, we noted that one such equivalence class was

$$\{134, 143, 314, 341, 413, 431\}.$$

The other three are

$$\{234, 243, 324, 342, 423, 432\},$$

$$\{132, 123, 312, 321, 213, 231\},$$

and

$$\{124, 142, 214, 241, 412, 421\}.$$

The product principle told us that if q is the number of such equivalence class, if each equivalence class has $k!$ elements, and the entire set of lists has $n!/(n - k)!$ element, then we must have that

$$qk! = n!/(n - k)! .$$

Dividing, we solve for q and get an expression for the number of k element subsets of an n element set. In fact, this is how we proved Theorem 1.2.

A principle that helps in learning and understanding mathematics is that if we have a mathematical result that shows a certain symmetry, it often helps our understanding to find a proof that reflects this symmetry. We call this the *Symmetry Principle*.

Principle 1.6 *If a formula has a symmetry (e.g. interchanging two variables doesn't change the result), then a proof that explains this symmetry is likely to give us additional insight into the formula.*

The proof above does not account for the symmetry of the $k!$ term and the $(n - k)!$ term in the expression $\frac{n!}{k!(n-k)!}$. This symmetry arises because choosing a k element subset is equivalent to choosing the $(n - k)$ -element subset of elements we don't want. In Exercise 1.4-4, we saw that the binomial coefficient $\binom{n}{k}$ also counts the number of ways to label n objects, say with the labels “in” and “out,” so that we have k “ins” and therefore $n - k$ “outs.” For each labelling, the k objects that get the label “in” are in our subset. This explains the symmetry in our formula, but it doesn't prove the formula. Here is a new proof that the number of labellings is $n!/k!(n - k)!$ that explains the symmetry.

Suppose we have m ways to assign k blue and $n - k$ red labels to n elements. From each labeling, we can create a number of lists, using the convention of listing the k blue elements first and the remaining $n - k$ red elements last. For example, suppose we are considering the number of ways to label 3 elements blue (and 2 red) from a five element set $\{A, B, C, D, E\}$. Consider

the particular labelling in which A , B , and D are labelled blue and C and E are labelled red. Which lists correspond to this labelling? They are

$$\begin{array}{cccccc} ABDCE & ABDEC & ADBCE & ADBEC & BADCE & BADEC \\ BDACE & BDAEC & DABCE & DABEC & DBACE & DBAEC \end{array}$$

that is, all lists in which A , B , and D precede C and E . Since there are $3!$ ways to arrange A , B , and D , and $2!$ ways to arrange C and E , by the product principle, there are $3!2! = 12$ lists in which A , B , and D precede C and E . For each of the q ways to construct a labelling, we could find a similar set of 12 lists that are associated with that labelling. Since *every* possible list of 5 elements will appear exactly once via this process, and since there are $5! = 120$ five-element lists overall, we must have by the product principle that

$$q \cdot 12 = 120, \tag{1.14}$$

or that $q = 10$. This agrees with our previous calculations of $\binom{5}{3} = 10$ for the number of ways to label 5 items so that 3 are blue and 2 are red.

Generalizing, we let q be the number of ways to label n objects with k blue labels and $n - k$ red labels. To create the lists associated with a labelling, we list the blue elements first and then the red elements. We can mix the k blue elements among themselves, and we can mix the $n - k$ red elements among themselves, giving us $k!(n - k)!$ lists consisting of first the elements with a blue label followed by the elements with a red label. Since we can choose to label any k elements blue, each of our lists of n distinct elements arises from some labelling in this way. Each such list arises from only one labelling, because two different labellings will have a different first k elements in any list that corresponds to the labelling. Each such list arises only once from a given labelling, because two different lists that correspond to the same labelling differ by a permutation of the first k places or the last $n - k$ places or both. Therefore, by the product principle, $qk!(n - k)!$ is the number of lists we can form with n distinct objects, and this must equal $n!$. This gives us

$$qk!(n - k)! = n!,$$

and division gives us our original formula for q . Recall that our proof of the formula we had in Exercise 1.4-5 did not explain why the product of three factorials appeared in the denominator, it simply proved the formula was correct. With this idea in hand, we could now explain *why* the product in the denominator of the formula in Exercise 1.4-5 for the number of labellings with three labels is what it is, and could generalize this formula to four or more labels.

Equivalence Relations

The process above divided the set of all $n!$ lists of n distinct elements into classes (another word for sets) of lists. In each class, all the lists are mutually equivalent, with respect to labeling with two labels. More precisely, two lists of the n objects are equivalent for defining labellings if we get one from the other by mixing the first k elements among themselves and mixing the last $n - k$ elements among themselves. Relating objects we want to count to sets of lists (so that each object corresponds to an set of equivalent lists) is a technique we can use to solve a wide variety of counting problems. (This is another example of abstraction.)

A relationship that divides a set up into mutually exclusive classes is called an **equivalence relation**.⁸ Thus, if

$$S = S_1 \cup S_2 \cup \dots \cup S_m$$

and $S_i \cap S_j = \emptyset$ for all i and j with $i \neq j$, then the relationship that says any two elements $x \in S$ and $y \in S$ are equivalent if and only if they lie in the same set S_i is an equivalence relation. The sets S_i are called *equivalence classes*, and, as we noted in Section 1.1 the family S_1, S_2, \dots, S_m is called a **partition** of S . One partition of the set $S = \{a, b, c, d, e, f, g\}$ is $\{a, c\}$, $\{d, g\}$, $\{b, e, f\}$. This partition corresponds to the following (boring) equivalence relation: a and c are equivalent, d and g are equivalent, and $b, e,$ and f are equivalent. A slightly less boring equivalence relation is that two letters are equivalent if typographically, their top and bottom are at the same height. This give the partition $\{a, c, e\}$, $\{b, d\}$, $\{f\}$, $\{g\}$.

Exercise 1.4-1 On the set of integers between 0 and 12 inclusive, define two integers to be related if they have the same remainder on division by 3. Which numbers are related to 0? to 1? to 2? to 3? to 4?. Is this relationship an equivalence relation?

In Exercise 1.4-1, the set of numbers related to 0 is the set $\{0, 3, 6, 9, 12\}$, the set to 1 is $\{1, 4, 7, 10\}$, the set related to 2 is $\{2, 5, 8, 11\}$, the set related to 3 is $\{0, 3, 6, 9, 12\}$, the set related to 4 is $\{1, 4, 7, 10\}$. A little more precisely, a number is related to one of 0, 3, 6, 9, or 12, if and only if it is in the set $\{0, 3, 6, 9, 12\}$, a number is related to 1, 4, 7, or 10 if and only if it is in the set $\{1, 4, 7, 10\}$ and a number is related to 2, 5, 8, or 11 if and only if it is in the set $\{2, 5, 8, 11\}$. Therefore the relationship is an equivalence relation.

The Quotient Principle

In Exercise 1.4-1 the equivalence classes had two different sizes. In the examples of counting labellings and subsets that we have seen so far, all the equivalence classes had the same size. This was very important. The principle we have been using to count subsets and labellings is given in the following theorem. We will call this principle the **Quotient Principle**.

Theorem 1.5 (Quotient Principle) *If an equivalence relation on a p -element set S has q classes each of size r , then $q = p/r$.*

Proof: By the product principle, $p = qr$, and so $q = p/r$. ■

Another statement of the quotient principle that uses the idea of a partition is

Principle 1.7 (Quotient Principle.) *If we can partition a set of size p into q blocks of size r , then $q = p/r$.*

Returning to our example of 3 blue and 2 red labels, $s = 5! = 120$, $t = 12$ and so by Theorem 1.5,

$$m = \frac{s}{t} = \frac{120}{12} = 10 .$$

⁸The usual mathematical approach to equivalence relations, which we shall discuss in the exercises, is different from the one given here. Typically, one sees an equivalence relation defined as a reflexive (everything is related to itself), symmetric (if x is related to y , then y is related to x), and transitive (if x is related to y and y is related to z , then x is related to z) relationship on a set X . Examples of such relationships are equality (on any set), similarity (on a set of triangles), and having the same birthday as (on a set of people). The two approaches are equivalent, and we haven't found a need for the details of the other approach in what we are doing in this course.

Equivalence class counting

We now give several examples of the use of Theorem 1.5.

Exercise 1.4-2 When four people sit down at a round table to play cards, two lists of their four names are equivalent as seating charts if each person has the same person to the right in both lists⁹. (The person to the right of the person in position 4 of the list is the person in position 1). We will use Theorem 1.5 to count the number of possible ways to seat the players. We will take our set S to be the set of all 4-element permutations of the four people, i.e., the set of all lists of the four people.

- (a) How many lists are equivalent to a given one?
- (b) What are the lists equivalent to ABCD?
- (c) Is the relationship of equivalence an equivalence relation?
- (d) Use the Quotient Principle to compute the number of equivalence classes, and hence, the number of possible ways to seat the players.

Exercise 1.4-3 We wish to count the number of ways to attach n distinct beads to the corners of a regular n -gon (or string them on a necklace). We say that two lists of the n beads are equivalent if each bead is adjacent to exactly the same beads in both lists. (The first bead in the list is considered to be adjacent to the last.)

- How does this exercise differ from the previous exercise?
- How many lists are in an equivalence class?
- How many equivalence classes are there?

In Exercise 1.4-2, suppose we have named the places at the table north, east, south, and west. Given a list we get an equivalent one in two steps. First we observe that we have four choices of people to sit in the north position. Then there is one person who can sit to this person's right, one who can be next on the right, and one who can be the following on on the right, all determined by the original list. Thus there are exactly four lists equivalent to a given one, including that given one. The lists equivalent to ABCD are ABCD, BCDA, CDAB, and DABC. This shows that two lists are equivalent if and only if we can get one from the other by moving everyone the same number of places to the right around the table (or we can get one from the other moving everyone the same number of places to the left around the table). From this we can see we have an equivalence relation, because each list is in one of these sets of four equivalent lists, and if two lists are equivalent, they are right or left shifts of each other, and we've just observed that all right and left shifts of a given list are in the same class. This means our relationship divides the set of all lists of the four names into equivalence classes each of size four. There are a total of $4! = 24$ lists of four distinct names, and so by Theorem 1.5 we have $4!/4 = 3! = 6$ seating arrangements.

Exercise 1.4-3 is similar in many ways to Exercise 1.4-2, but there is one significant difference. We can visualize the problem as one of dividing lists of n distinct beads up into equivalence classes,

⁹Think of the four places at the table as being called north, east, south, and west, or numbered 1-4. Then we get a list by starting with the person in the north position (position 1), then the person in the east position (position 2) and so on clockwise

but now two lists are equivalent if each bead is adjacent to exactly the same beads in both of them. Suppose we number the vertices of our polygon as 1 through n clockwise. Given a list, we can count the equivalent lists as follows. We have n choices for which bead to put in position 1. Then either of the two beads adjacent to it¹⁰ in the given list can go in position 2. But now, only one bead can go in position 3, because the other bead adjacent to position 2 is already in position 1. We can continue in this way to fill in the rest of the list. For example, with $n = 4$, the lists ABCD, ADCB, BCDA, BADC, CDAB, CBAD, DABC, and DCBA are all equivalent. Notice the first, third, fifth and seventh lists are obtained by shifting the beads around the polygon, as are the second, fourth, sixth and eighth (though in the opposite direction). Also note that the eighth list is the reversal of the first, the third is the reversal of the second, and so on. Rotating a necklace in space corresponds to shifting the letters in the list. Flipping a necklace over in space corresponds to reversing the order of a list. There will always be $2n$ lists we can get by shifting and reversing shifts of a list. The lists equivalent to a given one consist of everything we can get from the given list by rotations and reversals. Thus the relationship of every bead being adjacent to the same beads divides the set of lists of beads into disjoint sets. These sets, which have size $2n$, are the equivalence classes of our equivalence relation. Since there are $n!$ lists, Theorem 1.5 says there are

$$\frac{n!}{2n} = \frac{(n-1)!}{2}$$

bead arrangements.

Multisets

Sometimes when we think about choosing elements from a set, we want to be able to choose an element more than once. For example the set of letters of the word “roof” is $\{f, o, r\}$. However it is often more useful to think of the of the *multiset* of letters, which in this case is $\{\{f, o, o, r\}\}$. We use the double brackets to distinguish a multiset from a set. We can specify a *multiset* chosen from a set S by saying how many times each of its elements occurs. If S is the set of English letters, the “multiplicity” function for roof is given by $m(f) = 1$, $m(o) = 2$, $m(r) = 1$, and $m(\text{letter}) = 0$ for every other letter. In a multiset, order is not important, that is the multiset $\{\{r, o, f, o\}\}$ is equivalent to the multiset $\{\{f, o, o, r\}\}$. We know that this is the case, because they each have the same multiplicity function. We would like to say that the size of $\{\{f, o, o, r\}\}$ is 4, so we define the *size* of a multiset to be the sum of the multiplicities of its elements.

Exercise 1.4-4 Explain how placing k identical books onto the n shelves of a bookcase can be thought of as giving us a k -element multiset of the shelves of the bookcase. Explain how distributing k identical apples to n children can be thought of as giving us a k -element multiset of the children.

In Exercise 1.4-4 we can think of the multiplicity of a bookshelf as the number of books it gets and the multiplicity of a child as the number of apples the child gets. In fact, this idea of distribution of identical objects to distinct recipients gives a great mental model for a multiset chosen from a set S . Namely, to determine a k -element multiset chosen from S form S , we “distribute” k identical objects to the elements of S and the number of objects an element x gets is the multiplicity of x .

¹⁰Remember, the first and last bead are considered adjacent, so they have two beads adjacent to them.

Notice that it makes no sense to ask for the number of multisets we may choose from a set with n elements, because $\{\{A\}\}$, $\{\{A, A\}\}$, $\{\{A, A, A\}\}$, and so on are infinitely many multisets chosen from the set $\{A\}$. However it does make sense to ask for the number of k -element multisets we can choose from an n -element set. What strategy could we employ to figure out this number? To count k -element subsets, we first counted k -element permutations, and then divided by the number of different permutations of the same set. Here we need an analog of permutations that allows repeats. A natural idea is to consider lists with repeats. After all, one way to describe a multiset is to list it, and there could be many different orders for listing a multiset. However the two element multiset $\{\{A, A\}\}$ can be listed in just one way, while the two element multiset $\{\{A, B\}\}$ can be listed in two ways. When we counted k -element subsets of an n -element set by using the quotient principle, it was essential that each k -element subset corresponded to the same number (namely $k!$) of permutations (lists), because we were using the reasoning behind the quotient principle to do our counting here. So if we hope to use similar reasoning, we can't apply it to lists with repeats because different k -element multisets can correspond to different numbers of lists.

Suppose, however, we could count the number of ways to arrange k distinct books on the n shelves of a bookcase. We can still think of the multiplicity of a shelf as being the number of books on it. However, many different arrangements of distinct books will give us the same multiplicity function. In fact, any way of mixing the books up among themselves that does not change the number of books on each shelf will give us the same multiplicities. But the number of ways to mix the books up among themselves is the number of permutations of the books, namely $k!$. Thus it looks like we have an equivalence relation on the arrangements of distinct books on a bookshelf such that

1. Each equivalence class has $k!$ elements, and
2. There is a bijection between the equivalence classes and k -element multisets of the n shelves.

Thus if we can compute the number of ways to arrange k *distinct* books on the n shelves of a bookcase, we should be able to apply the quotient principle to compute the number of k -element multisets of an n -element set.

The bookcase arrangement problem.

Exercise 1.4-5 We have k books to arrange on the n shelves of a bookcase. The order in which the books appear on a shelf matters, and each shelf can hold all the books. We will assume that as the books are placed on the shelves they are moved as far to the left as they will go so that all that matters is the order in which the books appear and not the actual places where the books sit. When book i is placed on a shelf, it can go between two books already there or to the left or right of all the books on that shelf.

- (a) Since the books are distinct, we may think of a first, second, third, etc. book. In how many ways may we place the first book on the shelves?
- (b) Once the first book has been placed, in how many ways may the second book be placed?
- (c) Once the first two books have been placed, in how many ways may the third book be placed?

- (d) Once $i - 1$ books have been placed, book i can be placed on any of the shelves to the left of any of the books already there, but there are some additional ways in which it may be placed. In how many ways in total may book i be placed?
- (e) In how many ways may k distinct books be placed on n shelves in accordance with the constraints above?

Exercise 1.4-6 How many k -element multisets can we choose from an n -element set?

In Exercise 1.4-5 there are n places where the first book can go, namely on the left side of any shelf. Then the next book can go in any of the n places on the far left side of any shelf, or it can go to the right of book one. Thus there are $n + 1$ places where book 2 can go. At first, placing book three appears to be more complicated, because we could create two different patterns by placing the first two books. However book 3 could go to the far left of any shelf or to the immediate right of any of the books already there. (Notice that if book 2 and book 1 are on shelf 7 in that order, putting book 3 to the immediate right of book 2 means putting it between book 2 and book 1.) Thus in any case, there are $n + 2$ ways to place book 3. Similarly, once $i - 1$ books have been placed, there are $n + i - 1$ places where we can place book i . It can go at the far left of any of the n shelves or to the immediate right of any of the $i - 1$ books that we have already placed. Thus the number of ways to place k distinct books is

$$n(n+1)(n+2)\cdots(n+k-1) = \prod_{i=1}^k (n+i-1) = \prod_{j=0}^{k-1} (n+j) = \frac{(n+k-1)!}{(n-1)!}. \quad (1.15)$$

The specific product that arose in Equation 1.15 is called a *rising factorial power*. It has a notation (also introduced by Don Knuth) analogous to that for the falling factorial notation. Namely, we write

$$n^{\overline{k}} = n(n+1)\cdots(n+k-1) = \prod_{i=1}^k (n+i-1).$$

This is the product of k successive numbers beginning with n .

The number of k -element multisets of an n -element set.

We can apply the formula of Exercise 1.4-5 to solve Exercise 1.4-6. We define two bookcase arrangements of k books on n shelves to be equivalent if we get one from the other by permuting the books among themselves. Thus if two arrangements put the same number of books on each shelf they are put into the same class by this relationship. On the other hand, if two arrangements put a different number of books on at least one shelf, they are not equivalent, and therefore they are put into different classes by this relationship. Thus the classes into which this relationship divides the the arrangements are disjoint and partition the set of all arrangements. Each class has $k!$ arrangements in it. The set of all arrangements has $n^{\overline{k}}$ arrangements in it. This leads to the following theorem.

Theorem 1.6 *The number of k -element multisets chosen from an n -element set is*

$$\frac{n^{\overline{k}}}{k!} = \binom{n+k-1}{k}.$$

Proof: The relationship on bookcase arrangements that two arrangements are equivalent if and only if we get one from the other by permuting the books is an equivalence relation. The set of all arrangements has $n^{\overline{k}}$ elements, and the number of elements in an equivalence class is $k!$. By the quotient principle, the number of equivalence classes is $\frac{n^{\overline{k}}}{k!}$. There is a bijection between equivalence classes of bookcase arrangements with k books and multisets with k elements. The second equality follows from the definition of binomial coefficients. ■

The number of k -element multisets chosen from an n -elements is sometimes called the number of *combinations with repetitions* of n elements taken k at a time.

The right-hand side of the formula is a binomial coefficient, so it is natural to ask whether there is a way to interpret choosing a k -element *multiset* from an n -element set as choosing a k -element *subset* of some different $n + k - 1$ -element set. This illustrates an important principle. When we have a quantity that turns out to be equal to a binomial coefficient, it helps our understanding to interpret it as counting the number of ways to choose a subset of an appropriate size from a set of an appropriate size. We explore this idea for multisets in Problem 8 in this section.

Using the quotient principle to explain a quotient

Since the last expression in Equation 1.15 is quotient of two factorials it is natural to ask whether it is counting equivalence classes of an equivalence relation. If so, the set on which the relation is defined has size $(n + k - 1)!$. Thus it might be all lists or permutations of $n + k - 1$ distinct objects. The size of an equivalence class is $(n - 1)!$ and so what makes two lists equivalent might be permuting $n - 1$ of the objects among themselves. Said differently, the quotient principle suggests that we look for an explanation of the formula involving lists of $n + k - 1$ objects, of which $n - 1$ are identical, so that the remaining k elements are distinct. Can we find such an interpretation?

Exercise 1.4-7 In how many ways may we arrange k distinct books and $n - 1$ identical blocks of wood in a straight line?

Exercise 1.4-8 How does Exercise 1.4-7 relate to arranging books on the shelves of a bookcase?

In Exercise 1.4-7, if we tape numbers to the wood so that so that the pieces of wood are distinguishable, there are $(n + k - 1)!$ arrangements of the books and wood. But since the pieces of wood are actually indistinguishable, $(n - 1)!$ of these arrangements are equivalent. Thus by the quotient principle there are $(n + k - 1)! / (n - 1)!$ arrangements. Such an arrangement allows us to put the books on the shelves as follows: put all the books before the first piece of wood on shelf 1, all the books between the first and second on shelf 2, and so on until you put all the books after the last piece of wood on shelf n .

Important Concepts, Formulas, and Theorems

1. *Symmetry Principle.* If we have a mathematical result that shows a certain symmetry, it often helps our understanding to find a proof that reflects this symmetry.
2. *Partition.* Given a set S of items, a *partition* of S consists of m sets S_1, S_2, \dots, S_m , sometimes called *blocks* so that $S_1 \cup S_2 \cup \dots \cup S_m = S$ and for each i and j with $i \neq j$, $S_i \cap S_j = \emptyset$.

3. *Equivalence relation. Equivalence class.* A relationship that partitions a set up into mutually exclusive classes is called an **equivalence relation**. Thus if $S = S_1 \cup S_2 \cup \dots \cup S_m$ is a partition of S , the relationship that says any two elements $x \in S$ and $y \in S$ are equivalent if and only if they lie in the same set S_i is an equivalence relation. The sets S_i are called *equivalence classes*.
4. *Quotient principle.* The **quotient principle** says that if we can partition a set of p objects up into q classes of size r , then $q = p/r$. Equivalently, if an equivalence relation on a set of size p has q equivalence classes of size r , then $q = p/r$. The quotient principle is frequently used for counting the number of equivalence classes of an equivalence relation. When we have a quantity that is a quotient of two others, it is often helpful to our understanding to find a way to use the quotient principle to explain why we have this quotient.
5. *Multiset.* A multiset is similar to a set except that each item can appear multiple times. We can specify a *multiset* chosen from a set S by saying how many times each of its elements occurs.
6. *Choosing k -element multisets.* The number of k -element multisets that can be chosen from an n -element set is


$$\frac{(n+k-1)!}{k!(n-1)!} = \binom{n+k-1}{k}.$$

This is sometimes called the formula for “combinations with repetitions.”

7. *Interpreting binomial coefficients.* When we have a quantity that turns out to be a binomial coefficient (or some other formula we recognize) it is often helpful to our understanding to try to interpret the quantity as the result of choosing a subset of a set (or doing whatever the formula that we recognize counts.)

Problems

1. In how many ways may n people be seated around a round table? (Remember, two seating arrangements around a round table are equivalent if everyone is in the same position relative to everyone else in both arrangements.)
2. In how many ways may we embroider n circles of different colors in a row (lengthwise, equally spaced, and centered halfway between the top and bottom edges) on a scarf (as follows)?



3. Use binomial coefficients to determine in how many ways three identical red apples and two identical golden apples may be lined up in a line. Use equivalence class counting (in particular, the quotient principle) to determine the same number.

4. Use multisets to determine the number of ways to pass out k identical apples to n children.
5. In how many ways may n men and n women be seated around a table alternating gender? (Use equivalence class counting!!)
6. In how many ways may we pass out k identical apples to n children if each child must get at least one apple?
7. In how many ways may we place k distinct books on n shelves of a bookcase (all books pushed to the left as far as possible) if there must be at least one book on each shelf?
8. The formula for the number of multisets is $(n + k - 1)!$ divided by a product of two other factorials. We seek an explanation using the quotient principle of why this counts multisets. The formula for the number of multisets is also a binomial coefficient, so it should have an interpretation involving choosing k items from $n + k - 1$ items. The parts of the problem that follow lead us to these explanations.
 - (a) In how many ways may we place k red checkers and $n - 1$ black checkers in a row?
 - (b) How can we relate the number of ways of placing k red checkers and $n - 1$ black checkers in a row to the number of k -element multisets of an n -element set, say the set $\{1, 2, \dots, n\}$ to be specific?
 - (c) How can we relate the choice of k items out of $n + k - 1$ items to the placement of red and black checkers as in the previous parts of this problem?
9. How many solutions to the equation $x_1 + x_2 + \dots + x_n = k$ are there with each $x_i \geq 0$?
10. How many solutions to the equation $x_1 + x_2 + \dots + x_n = k$ are there with each $x_i > 0$?
11. In how many ways may n red checkers and $n + 1$ black checkers be arranged in a circle? (This number is a famous number called a *Catalan number*.)
12. A standard notation for the number of partitions of an n element set into k classes is $S(n, k)$. $S(0, 0)$ is 1, because technically the empty family of subsets of the empty set is a partition of the empty set, and $S(n, 0)$ is 0 for $n > 0$, because there are no partitions of a nonempty set into no parts. $S(1, 1)$ is 1.
 - (a) Explain why $S(n, n)$ is 1 for all $n > 0$. Explain why $S(n, 1)$ is 1 for all $n > 0$.
 - (b) Explain why, for $1 < k < n$, $S(n, k) = S(n - 1, k - 1) + kS(n - 1, k)$.
 - (c) Make a table like our first table of binomial coefficients that shows the values of $S(n, k)$ for values of n and k ranging from 1 to 6.
13. You are given a square, which can be rotated 90 degrees at a time (i.e. the square has four orientations). You are also given two red checkers and two black checkers, and you will place each checker on one corner of the square. How many lists of four letters, two of which are R and two of which are B, are there? Once you choose a starting place on the square, each list represents placing checkers on the square in clockwise order. Consider two lists to be equivalent if they represent the same arrangement of checkers at the corners of the square, that is, if one arrangement can be rotated to create the other one. Write down the equivalence classes of this equivalence relation. Why can't we apply Theorem 1.5 to compute the number of equivalence classes?

14. The terms “reflexive”, “symmetric” and “transitive” were defined in Footnote 2. Which of these properties is satisfied by the relationship of “greater than?” Which of these properties is satisfied by the relationship of “is a brother of?” Which of these properties is satisfied by “is a sibling of?” (You are not considered to be your own brother or your own sibling). How about the relationship “is either a sibling of or is?”
- Explain why an equivalence relation (as we have defined it) is a reflexive, symmetric, and transitive relationship.
 - Suppose we have a reflexive, symmetric, and transitive relationship defined on a set S . For each x in S , let $S_x = \{y | y \text{ is related to } x\}$. Show that two such sets S_x and S_y are either disjoint or identical. Explain why this means that our relationship is an equivalence relation (as defined in this section of the notes, not as defined in the footnote).
 - Parts b and c of this problem prove that a relationship is an equivalence relation if and only if it is symmetric, reflexive, and transitive. Explain why. (A short answer is most appropriate here.)
15. Consider the following C++ function to compute $\binom{n}{k}$.

```
int pascal(int n, int k)
{
    if (n < k)
    {
        cout << "error: n<k" << endl;
        exit(1);
    }

    if ( (k==0) || (n==k) )
        return 1;

    return pascal(n-1,k-1) + pascal(n-1,k);
}
```

Enter this code and compile and run it (you will need to create a simple main program that calls it). Run it on larger and larger values of n and k , and observe the running time of the program. It should be surprisingly slow. (Try computing, for example, $\binom{30}{15}$.) Why is it so slow? Can you write a different function to compute $\binom{n}{k}$ that is *significantly faster*? Why is your new version faster? (Note: an exact analysis of this might be difficult at this point in the course; it will be easier later. However, you should be able to figure out roughly why this version is so much slower.)

16. Answer each of the following questions with either n^k , $n^{\underline{k}}$, $\binom{n}{k}$, or $\binom{n+k-1}{k}$.
- In how many ways can k different candy bars be distributed to n people (with any person allowed to receive more than one bar)?
 - In how many ways can k different candy bars be distributed to n people (with nobody receiving more than one bar)?

- (c) In how many ways can k identical candy bars distributed to n people (with any person allowed to receive more than one bar)?
- (d) In how many ways can k identical candy bars distributed to n people (with nobody receiving more than one bar)?
- (e) How many one-to-one functions f are there from $\{1, 2, \dots, k\}$ to $\{1, 2, \dots, n\}$?
- (f) How many functions f are there from $\{1, 2, \dots, k\}$ to $\{1, 2, \dots, n\}$?
- (g) In how many ways can one choose a k -element subset from an n -element set?
- (h) How many k -element multisets can be formed from an n -element set?
- (i) In how many ways can the top k ranking officials in the US government be chosen from a group of n people?
- (j) In how many ways can k pieces of candy (not necessarily of different types) be chosen from among n different types?
- (k) In how many ways can k children each choose one piece of candy (all of different types) from among n different types of candy?