

E0234: Assignment 5

Due: Monday, 15th Feb 2016.

It is highly recommended you do not google for the answers to the questions below. You can discuss with your friends, but then mention that in your submission. The writing should solely be your own.

- Let X_1, \dots, X_n be independent indicator random variables, and let $p_i := \Pr[X_i = 1]$. Let U and L be any positive reals such that $U \geq \sum_{i=1}^n p_i \geq L$. Let $X = \sum_{i=1}^n X_i$. Prove
 - $\Pr[X \geq (1 + \delta)U] \leq e^{-U((1+\delta)\ln(1+\delta)-\delta)}$
 - $\Pr[X \leq (1 - \delta)L] \leq e^{-L\delta^2/2}$
- We are given a collection of n sets S_1, \dots, S_n where each S_i is a subset of $U = \{1, 2, \dots, n\}$. A 2-coloring is an assignment $c : U \rightarrow \{-1, +1\}$ to the elements of U . The *discrepancy* of a 2-coloring is $\max_{j=1}^n \sum_{i \in S_j} c(i)$. What can you say about the discrepancy of a random coloring which independently assigns each element $+1$ or -1 with probability $1/2$? In particular, give a confidence bound (in terms of n) in which the discrepancy lies with probability $> 99\%$.
- If Z is a random variable with $\mathbf{Exp}[Z] = 0$ and $|Z| \leq 1$ with probability 1, then prove Z is 1-subgaussian.
 - If Z is a σ -subgaussian random variable, prove that $\mathbf{Exp}[Z] = 0$ and $\mathbf{Var}[Z] \leq \sigma^2$.
- Johnson-Lindenstrauss Lower Bound.** Consider the following $(n+1)$ points in n dimensions: $\{0, e_1, \dots, e_n\}$ where e_i is the i th unit vector with 1 in the i th position and 0 elsewhere. Consider a mapping of these $(n+1)$ vectors to k dimensions such that the pairwise distances are preserved to a $(1 \pm \varepsilon)$ -multiplicative factor. What is the best *lower bound* you can prove on k ?